

## HERBICIDE

# BADGE® II

Get proven control of almost 30 broadleaf weeds in cereals, flax and corn with easy-to-use tank-mix options for one-shot weed control.



### ACTIVE INGREDIENTS

Bromoxynil and MCPA ester

### CHEMISTRY GROUPS

Group 4 (MCPA ester) and Group 6 (bromoxynil)

### APPLICATION RATES AND PACKAGING

- 0.5 L/ac or 1 case treats 40 acres
- 2 x 10 L jug/case

### REGISTERED CROPS:

#### Field crops:

- Barley
- Canary seed
- Corn
- Fall rye

- Flax
- Oats
- Solin

- Timothy (established for seed production)
- Wheat (spring, winter, durum)

#### Seedling grasses:

- Bromegrass
- Creeping red fescue
- Crested wheatgrass
- Intermediate wheatgrass
- Meadow fescue

- Meadow foxtail
- Orchard grass
- Reed canarygrass
- Russian wild rye
- Seedling meadow bromegrass

- Seedling tall fescue
- Slender wheatgrass
- Streambank wheatgrass
- Tall wheatgrass
- Timothy

### WEEDS CONTROLLED:

- American nightshade
- Ball mustard
- Bluebur
- Canada thistle<sup>1</sup>
- Cocklebur
- Common buckwheat
- Common groundsel
- Common ragweed
- Cow cockle
- Flixweed
- Green smartweed
- Kochia<sup>2</sup>

- Lady's thumb
- Lamb's quarters
- Night-flowering catchfly
- Pale smartweed
- Perennial sow thistle<sup>1</sup>
- Redroot pigweed
- Russian thistle<sup>2</sup>
- Scentless chamomile<sup>3</sup>
- Shepherd's purse
- Stinkweed

- Tartary buckwheat
- Velvetleaf<sup>4</sup>
- Volunteer canola (all types)
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild tomato
- Wormseed mustard

<sup>1</sup>Top growth control.

<sup>2</sup>Spray before plants are 2 inches high.

<sup>3</sup>Spring annual only.

<sup>4</sup>Spray before plants are 3 inches high.

<sup>5</sup>Up to 4-leaf stage.

### HOW IT WORKS:

A combination of systemic and contact activity with weeds yellowing within 2–4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2–10 days.

**BADGE® II****CROP STAGING:**

Crop	Stage
Barley, Oats, Spring wheat (including durum)	2 leaf to early flag
Canary seed	3 – 5 leaf
Corn	4 – 6 leaf
Fall rye	When growth commences in spring to early flag leaf.
Flax, Solin	2 inches to early bud stage. Best tolerance occurs when flax is 2 – 4 inches tall.
Seedling grasses	2 – 4 leaf
Timothy (established for seed production)	Prior to shot blade in the seed production year.
Winter wheat	2 – 4 leaf stage in the fall or after growth resumes up to early flag leaf.

**WATER VOLUME:**

- Ground: 20 – 40 L/ac in cereals and flax; 80 – 120 L/ac in corn; 60 L/ac in forages
- Aerial: 8 – 20 L/ac

**RAINFASTNESS:**

1 hour

**SUPPORTED TANK MIXES:**

- Corn: Atrazine
- Flax and Solin: Clethodim (Arrow® 240 EC) or Poast®
- Oats: MCPA
- Spring wheat and barley: Ally®, Avenge®, MCPA, Refine Extra® or tralkoxydim (Bison® 400 L)
- Winter wheat: Refine Extra

**MIXING INSTRUCTIONS:**

1. Fill clean spray tank ½ full with water.
2. Add the required amount of Badge® II and agitate thoroughly.
3. Fill the tank and agitate again before use.
4. When tank mixing, follow instructions on both labels.

**CROP ROTATIONS:**

No re-cropping restrictions the year after treatment.

**GRAZING RESTRICTIONS:**

- Do not graze treated grain or established timothy crops or cut for feed within 30 days of application.
- Do not graze meadow foxtail in the year of treatment.
- Do not graze other treated forage grasses within 56 days of treatment.

**PRE-HARVEST INTERVAL:**

Flax or Solin: 60 days

**STORAGE:**

Avoid freezing.

**QUICK TIPS:**

Badge II herbicide is well known for being gentle on the crop. Avoid spraying if temperatures are above 25 C.