

HERBICIDE

BADGE® II

Get proven control of almost 30 broadleaf weeds in cereals, flax and corn with easy-to-use tank-mix options for one-shot weed control.



ACTIVE INGREDIENTS

Bromoxynil and MCPA ester

CHEMISTRY GROUPS

Group 4 (MCPA ester) and Group 6 (bromoxynil)

APPLICATION RATES AND PACKAGING

- 0.5 L/ac or 1 case treats 40 acres
- 2 x 10 L jug/case
- 120 L drum treats 240 acres

REGISTERED CROPS:

Field crops:

- Barley
- Canary seed
- Corn
- Fall rye

- Flax
- Oats
- Solin

- Timothy (established for seed production)
- Wheat (spring, winter, durum)

Seedling grasses:

- Bromegrass
- Creeping red fescue
- Crested wheatgrass
- Intermediate wheatgrass
- Meadow fescue

- Meadow foxtail
- Orchard grass
- Reed canarygrass
- Russian wild rye
- Seedling meadow bromegrass

- Seedling tall fescue
- Slender wheatgrass
- Streambank wheatgrass
- Tall wheatgrass
- Timothy

WEEDS CONTROLLED:

- American nightshade
- Ball mustard
- Bluebur
- Canada thistle¹
- Cocklebur
- Common buckwheat
- Common groundsel
- Common ragweed
- Cow cockle⁵
- Flixweed
- Green smartweed
- Kochia²

- Lady's thumb
- Lamb's quarters
- Night-flowering catchfly
- Pale smartweed
- Perennial sow thistle¹
- Redroot pigweed
- Russian thistle²
- Scentless chamomile³
- Shepherd's purse
- Stinkweed

- Tartary buckwheat
- Velvetleaf⁴
- Volunteer canola (all types)
- Volunteer sunflower
- Wild buckwheat
- Wild mustard
- Wild tomato
- Wormseed mustard

¹Top growth control.

²Spray before plants are 2 inches high.

³Spring annual only.

⁴Spray before plants are 3 inches high.

⁵Up to 4-leaf stage.

HOW IT WORKS:

A combination of systemic and contact activity with weeds yellowing within 2–4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2–10 days.

BADGE® II**CROP STAGING:**

Crop	Stage
Barley, Oats, Spring wheat (including durum)	2 leaf to early flag
Canary seed	3 – 5 leaf
Corn	4 – 6 leaf
Fall rye	When growth commences in spring to early flag leaf.
Flax, Solin	2 inches to early bud stage. Best tolerance occurs when flax is 2 – 4 inches tall.
Seedling grasses	2 – 4 leaf
Timothy (established for seed production)	Prior to shot blade in the seed production year.
Winter wheat	2 – 4 leaf stage in the fall or after growth resumes up to early flag leaf.

WATER VOLUME:

- Ground: 20 – 40 L/ac in cereals and flax; 80 – 120 L/ac in corn; 60 L/ac in forages
- Aerial: 8 – 20 L/ac

RAINFASTNESS:

1 hour

SUPPORTED TANK MIXES:

- Corn: Atrazine
- Flax and Solin: Clethodim (Arrow® 240 EC) or Poast®
- Oats: MCPA
- Spring wheat and barley: Ally®, Avenge®, MCPA, Refine® SG or tralkoxydim (Bison® 400 L)
- Spring wheat only: Axial®, clodinafop (Ladder® 240 EC, Ladder® All-In) or Everest®
- Winter wheat: Refine Extra®

MIXING INSTRUCTIONS:

1. Fill clean spray tank ½ full with water.
2. Add the required amount of Badge® II and agitate thoroughly.
3. Fill the tank and agitate again before use.
4. When tank mixing, follow instructions on both labels.

CROP ROTATIONS:

No re-cropping restrictions the year after treatment.

GRAZING RESTRICTIONS:

- Do not graze treated grain or established timothy crops or cut for feed within 30 days of application.
- Do not graze meadow foxtail in the year of treatment.
- Do not graze other treated forage grasses within 56 days of treatment.

PRE-HARVEST INTERVAL:

Flax or Solin: 60 days

STORAGE:

Avoid freezing.

QUICK TIPS:

Badge II herbicide is well known for being gentle on the crop. Avoid spraying if temperatures are above 25 C.