

HERBICIDE

# BROMOTRIL® II POST-EMERGENT

Get post-emergent control of tough broadleaf weeds with tank-mix flexibility and excellent crop safety in cereals and other crops.



**ACTIVE INGREDIENT**

Bromoxynil

**CHEMISTRY GROUP**

Group 6

**APPLICATION RATES AND PACKAGING**

- 0.49 – 0.57 L/ac or 40 – 34 ac/case
- 2 x 9.7 L jugs/case

**REGISTERED CROPS:**

Crop	Crop Leaf Stage
Alfalfa (seedling)	2 – 6 trifoliolate
Alfalfa (established)	Spring: before the crop begins to shield the weeds
Barley, Oats, Triticale, Wheat (spring, durum)	2 leaf to early flag
Winter wheat	Fall: 2 – 4 leaf Spring: first growth to early flag
Corn (field, sweet) with drop pipes	Beyond 8 leaf
Canary seed (seed production only)	3 – 5 leaf
Fall rye	Spring: from first growth to early flag
Flax	2 – 4 inches in height
Forage millet, Sorghum	4 leaf to 8 inches
Seedling grasses (seed production only): Bromegrass, Fescue (creeping red, meadow), Orchard grass, Reed canarygrass, Russian wildrye, Timothy, Wheatgrass (crested, intermediate, slender, tall)	2 – 4 leaf (year of establishment only)

**WEEDS CONTROLLED:**

Seedling up to 4-leaf stage:

- American nightshade
- Bluebur
- Cocklebur
- Common ragweed
- Cow cockle<sup>1</sup>
- Green smartweed
- Kochia<sup>2</sup>
- Lady's thumb
- Pale smartweed
- Pigweed<sup>1</sup>
- Russian thistle<sup>2</sup>
- Stinkweed<sup>1</sup>
- Velvetleaf<sup>3</sup>
- Wild mustard<sup>1</sup>

Seedling up to 8-leaf stage:

- Common buckwheat
- Common groundsel
- Lamb's quarters
- Tartary buckwheat
- Wild buckwheat

<sup>1</sup>In normal conditions, it will be controlled up to 4-leaf stage. Plants beyond this stage are unlikely to be controlled; the higher rate generally gives better results.

<sup>2</sup>Spray before plants are 2 inches high.

<sup>3</sup>Spray before plants are 3 inches high.

**BROMOTRIL® II** POST-EMERGENT**HOW IT WORKS:**

Bromotrill® II is a contact herbicide. Leaves will yellow in 2–4 days with complete control in 7–14 days.

**WATER VOLUME:**

- Ground: 20–40 L/ac
- Air (wheat and barley only): 8–16 L/ac

**RAINFASTNESS:**

30 minutes

**SUPPORTED TANK MIXES:**

- **Spring wheat:** 2,4-D, Avenge®, MCPA, tralkoxydim (Bison® 400 L)
- **Winter wheat:** 2,4-D, MCPA, tralkoxydim (Bison 400 L)
- **Barley:** 2,4-D, Avenge, MCPA, tralkoxydim (Bison 400 L)
- **Oats:** MCPA
- **Corn:** Accent®, Atrazine, Banvel®, Ultim®
- **Fall rye:** MCPA
- **Flax:** MCPA or MCPA-K
- **Canary seed:** MCPA
- **Seedling grasses:** MCPA

**MIXING INSTRUCTIONS:**

1. Fill spray tank ½ full with water.
2. Add required amount of Bromotrill II. Begin agitation.
3. If tank mixing, add the recommended amount of the tank-mix partner product to the spray tank first, agitate and then add Bromotrill II (unless otherwise directed by the Bromotrill II and tank-mix partner label).
4. Add the remaining amount of water while agitation continues.

**CROP ROTATIONS:**

No restrictions.

**PRE-HARVEST INTERVAL:**

30 days

**GRAZING RESTRICTIONS:**

- Do not use treated crops for grazing of livestock or green feed until 30 days after application.
- Do not cut treated crops for forage until 30 days after application.

**STORAGE:**

Avoid freezing.

**QUICK TIPS:**

Avoid spraying if temperatures are above 25 C. Leaf scorching may occur in corn and flax if applied during or after adverse growing conditions, such as cool and wet or hot (above 27 C) and humid weather. For best results, spray when weeds are in the seedling stage.