

HERBICIDE

BROMOTRIL® II PRE-SEED

When tank mixed with glyphosate, Bromotril® II provides outstanding burn-off control of tough weeds like kochia, buckwheat, cow cockle and Russian thistle.



ACTIVE INGREDIENT

Bromoxynil

CHEMISTRY GROUP

Group 6

APPLICATION RATES AND PACKAGING

- Bromotril II at 0.49 L/ac (40 ac/case) tank mixed with glyphosate at a rate of 175 g a.i./ac
- 2 x 9.7 L jugs/case
- 116.4 L drum treats 240 acres

REGISTERED CROPS:

Tank mixed with glyphosate, this pre-seed burn-off should only be used for minimum or zero-tillage cropping systems in wheat, barley and oats.

WEEDS CONTROLLED:

Seedling up to 4-leaf stage:

- American nightshade
- Bluebur
- Cocklebur
- Common ragweed
- Cow cockle¹
- Green smartweed
- Kochia²
- Lady's thumb
- Pale smartweed
- Pigweed¹
- Russian thistle²
- Stinkweed¹
- Velvetleaf³
- Wild mustard¹

Seedling up to 8-leaf stage:

- Common buckwheat
- Common groundsel
- Lamb's quarters
- Tartary buckwheat
- Wild buckwheat

Grassy weeds up to 6 inches:

- Green foxtail
- Volunteer cereals
- Wild oats

¹In normal conditions, it will be controlled up to 4-leaf stage. Plants beyond this stage are unlikely to be controlled; the higher rate generally gives better results.

²Spray before plants are 2 inches high. Includes Group 2 and Group 9 resistant biotypes.

³Spray before plants are 3 inches high.

HOW IT WORKS:

Bromotril II is a contact herbicide. Leaves will yellow in 2–4 days with complete control in 7–14 days. Bromotril II controls Group 2 and Group 9 (glyphosate) resistant biotypes.

CROP STAGING:

- Application can be made just prior to or immediately after seeding.
- **Under no circumstances should application be made after crop emergence.**

WATER VOLUME:

Ground: 20–40 L/ac

HERBICIDE

BROMOTRIL® II PRE-SEED

RAINFASTNESS:

30 minutes

SUPPORTED TANK MIXES:

Must be tank mixed with glyphosate at a rate of 175 g a.i./ac of glyphosate.

MIXING INSTRUCTIONS:

1. Fill spray tank ½ full with water.
2. Add required amount of Bromotrill® II. Begin agitation.
3. Add the recommended amount of glyphosate to the spray tank, agitate.
4. Add surfactant.
5. Add the remaining amount of water while agitation continues.

CROP ROTATIONS:

Only apply prior to wheat, barley and oats.

STORAGE:

Avoid freezing.

QUICK TIPS:

For best results, spray when weeds are in the seedling stage.