

FUNGICIDE

FOLPAN® 80 WDG

A water-dispersible granular fungicide that delivers exceptional protection against the most common and economically damaging diseases in grapes and other fruit, vegetable and ornamental crops.

ACTIVE INGREDIENT

Folpet

CHEMISTRY GROUP

Group M

APPLICATION RATES AND PACKAGING

- 0.5–2 kg/ac
- 0.5 kg/ac–20 ac/pack
- 2 kg/ac–5 ac/pack
- 10 kg pack

REGISTERED CROPS:

- Apples
- Azalea
- Carnation
- Chrysanthemum
- Crabapples
- Field cucumbers
- Field tomatoes
- Grapes
- Iris
- Marigold
- Melons
- Poinsettia (greenhouse)
- Pumpkins
- Snapdragon
- Squash
- Strawberries
- Zinnia

KEY DISEASES CONTROLLED:

- Alternaria leaf spot
- Anthracnose
- Black rot
- Brooks spot
- Dead arm
- Didymellina leaf spot
- Downy mildew
- Fly-speck
- Fruit rot
- Gray mold
- Leaf spot
- Phythium root rot
- Powdery mildew
- Scab
- Septoria leaf spot
- Sooty blotch
- Stem rot

HOW IT WORKS:

Multi-site-contact mode of action with protective ability. Use in a regularly scheduled maintenance program.

APPLICATION TIMING AND CROP STAGING:

Crop	Diseases	Timing
Apples, Crabapples	Alternaria leaf spot, Black rot, Brooks spot, Fly-speck, Scab, Sooty blotch	Apply in a regular protective schedule from green tip up until harvest. Do not apply more than 6 applications per season. See label for additional precautions.
Cranberries	Fruit rot	Apply when 5% of blossoms are open. Repeat 10–14 days later when 50–75% of blossoms are open.
Grapes	Dead arm	Apply when new shoots are 1–2 inches long and repeat when growth is 4–6 inches long.
	Black rot, Downy mildew	Apply just before bloom, just after bloom and in first cover spray. For downy mildew, an additional application 2–3 weeks later may be needed.
	Powdery mildew	Apply in a tank mix with Kumulus® DF at first sign of mildew and repeat after 10 days.

FOLPAN® 80 WDG

APPLICATION TIMING AND CROP STAGING (continued):

Crop	Diseases	Timing
Field cucumbers, Melons, Pumpkins, Squash	Anthracnose, Downy mildew	Apply when first true leaves appear. Repeat at 7-day intervals until crop is harvested.
Field tomatoes	Anthracnose	Apply during first bloom and repeat at 7-day intervals until harvest.
Strawberries	Gray mold, Fruit rot, Leaf spot	Apply before first infection and repeat at 7-day intervals to protect crop until harvest.
Carnation, Poinsettia, Azalea, Marigold, Zinnia, Chrysanthemum, Iris, Snapdragon	Blight, Phythium root rot, Stem rot, Alternaria leaf spot, Septoria leaf spot, Didymellina leaf spot, Anthracnose (depending on ornamental)	Generally, apply when ornamental emerges and repeat at regular intervals. Consult label for timing on specific disease and ornamental.

WATER VOLUME:

Ground application only:
400–1200 L/ac

RAINFASTNESS:

No restrictions.

SUPPORTED TANK MIXES:

- Most commonly used insecticides, adjuvants and fungicides. Check compatibilities before making tank mixes.
- Fungicidal activity is reduced if combined with strongly alkaline materials such as hydrated lime.
- Do not use in combination with or closely following an oil spray.

MIXING INSTRUCTIONS:

1. Fill spray tank nearly full and pour recommended amount of Folpan® 80 WDG on surface of water.
2. Fungicide can be premixed in a bucket ½ filled with water. Mix can be poured through screen into nearly filled spray tank.
3. Finish filling tank.
4. Keep agitator running during filling and spraying.

CROP ROTATIONS:

No restrictions.

PRE-HARVEST INTERVALS:

- Cranberries: 30 days
- Field cucumbers: 1 day
- Field tomatoes: 1 day
- Grapes: 1 day
- Melons: 1 day
- Pumpkins: 1 day
- Squash: 1 day
- Strawberries: 1 day

GRAZING RESTRICTIONS:

No restrictions.

STORAGE:

No restrictions.

QUICK TIPS:

Folpan 80 WDG can be combined with most commonly used insecticides, adjuvants and fungicides. Check compatibility before tank mixing.