

HERBICIDE

FORCEFIGHTER™ M

It has 3 active ingredients and 2 modes of action to fight herbicide resistance and provide post-emergent control of broadleaf weeds in wheat and barley.



ACTIVE INGREDIENTS

Fluroxypyr, bromoxynil and MCPA ester

CHEMISTRY GROUPS

Group 4 (MCPA ester, fluroxypyr) and Group 6 (bromoxynil)

APPLICATION RATES AND PACKAGING

- 2 x 10 L bromoxynil/MCPA + 9.6 L fluroxypyr or 40 ac/case
- 2 x 120 L bromoxynil/MCPA + 115.2 L fluroxypyr treats 480 acres

REGISTERED CROPS:

- Wheat (spring, winter, durum)
- Barley

WEEDS CONTROLLED:

- American nightshade
- Bluebur
- Burdock
- Canada thistle¹
- Chickweed
- Cleavers (including Group 2 resistant biotypes)
- Cockelbur
- Common groundsel
- Cow cockle
- Flixweed
- Hempnettle
- Kochia (including Group 2 and glyphosate-resistant biotypes)
- Lady's thumb
- Lamb's quarters
- Night-flowering catchfly
- Perennial sow thistle
- Redroot pigweed
- Russian thistle
- Shepherd's purse
- Smartweed
- Stinkweed
- Vetch
- Volunteer canola
- Volunteer sunflower
- Wild buckwheat
- Wild mustard (including Group 2 resistant biotypes)
- Wild radish
- Wormseed mustard

¹Top growth control

HOW IT WORKS:

Quickly causes plants to stop growing. Convenient co-pack controls a wide range of weeds, including glyphosate-resistant and Group 2 resistant kochia, Group 2 resistant cleavers and Group 2 resistant wild mustard.

CROP STAGING:

Crop	Stage
Wheat	2 leaf to early flag
Barley	2 leaf to early flag

WATER VOLUME:

Ground: 20 – 40 L/ac

RAINFASTNESS:

1 hour

FORCEFIGHTER™ M

SUPPORTED TANK MIXES:

Wheat:

- Tralkoxydim (Bison® 400 L)
- Clodinafop (Ladder® 240 EC, Ladder® All-In)
- Simplicity®
- Thifensulfuron/tribenuron for suppression of narrow-leaved hawk's beard
- Refine® SG

Durum:

- Clodinafop (Ladder 240 EC, Ladder® All-In)
- Simplicity

Barley:

- Tralkoxydim (Bison 400 L)
- Thifensulfuron/tribenuron for suppression of narrow-leaved hawk's beard

MIXING INSTRUCTIONS:

1. Fill spray tank ½ full with water.
2. Add the required amount of Forcefighter™ M and agitate thoroughly.
3. Add any tank-mix partners.
4. Fill the tank and agitate again before use.

CROP ROTATIONS:

Can be seeded the following year to barley, canola, flax, forage grasses, lentils, mustard, peas, rye and wheat.

PRE-HARVEST INTERVAL:

60 days

GRAZING RESTRICTIONS:

30 days

STORAGE:

Store in a heated facility. If product is frozen, bring to room temperature and agitate well before use.

QUICK TIPS:

Forcefighter M's activity is influenced by weather conditions. The optimal temperature is 12–24 C. Avoid application 3 days before or after frost. Do not apply before the 2-leaf stage as crop injury may occur. Use 10 gal/ac application volume when there is a heavy canopy or when most weeds are at an advanced stage of growth. Do not apply by air.