

2020-0686
2020-11-06

GROUP

2

HERBICIDE

NEXT 70 WDG HERBICIDE

Water Dispersible Granular Herbicide

in water-soluble bags

COMMERCIAL
(AGRICULTURAL)

ACTIVE INGREDIENT: Imazamox 70% a.e.

REGISTRATION NO. 33620 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 50 g – 20 kg (117.5 g water-soluble bags)

WARNING - EYE IRRITANT

READ THE LABEL AND THIS BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

DO NOT SELL INDIVIDUAL WATER-SOLUBLE BAGS SEPARATELY

Sharda Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
Sharda Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, Canada
M9R1V1
1-844-810-5720
1-416-840-5639

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT APPLY using aerial application equipment.

Skin irritant.

WARNING - EYE IRRITANT

May cause eye damage.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, goggles or face shield, socks and shoes during mixing, loading, clean-up and repair. Goggle or faceshield are not required during application and gloves are not required during application within a closed cab.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

Store above 5°C.

Store in a cool, dry place away from children, animals, food, feed or fertilizers.

Keep unused water-soluble bags in resealed original container.
Keep packages dry at all times.
Keep packages out of rain.
Keep from freezing.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
Avoid application of this product when heavy rain is forecast.
Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

For use only on on soybeans in Eastern Canada only and field peas, soybeans imazamox tolerant Canola imazamox tolerant Lentils imazamox tolerant Sunflowers and imazamox tolerant Canola Quality *Brassica juncea* in the Prairie Provinces and Interior of British Columbia (including the Peace River Region) only.

DIRECTIONS FOR USE (Eastern Canada Only)

NEXT 70 WDG Herbicide is a selective herbicide that can be applied as an early post-mergence treatment (cotyledon to 4-leaf stage) in soybeans.

DIRECTIONS FOR USE (Prairie Provinces and Interior of British Columbia (including the Peace River Region only)

NEXT 70 WDG Herbicide is a selective herbicide that can be applied as an early post-emergence treatment in field peas, soybeans, imazamox tolerant Canola, imazamox tolerant Lentils, imazamox tolerant Sunflowers and imazamox tolerant Canola Quality *Brassica juncea*.

When NEXT 70 WDG Herbicide is applied early post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

DO NOT apply more than once per year.

DO NOT graze the treated field peas, soybeans, imazamox tolerant Canola, imazamox tolerant Lentils or imazamox tolerant Canola quality *Brassica juncea* or cut for hay within 20 days of application.

DO NOT graze treated sunflower plants or cut for straw; sufficient data are not available to support such use.

Pre-harvest Interval: After 60 days, dry edible beans (*Phaseolus vulgaris*), field peas, soybeans, canola, imazamox tolerant Lentils and imazamox tolerant Canola Quality *Brassica juncea* can be harvested. Sunflowers can be harvested after 70 days.

Field Sprayer Application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply using aerial application equipment.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

Buffer Zones

Spot treatments using hand-held equipment DO NOT require a buffer zone.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands).

| Method of Application | Crop | Buffer zones (metres) Required for the Protection of Terrestrial Habitat: |
|-----------------------|-----------|---|
| Field sprayer | All crops | 1 |

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

BROADLEAF WEED AND GRASS CONTROL (Prairie Provinces and Interior of British Columbia (including the Peace River Region) only)

| | |
|--|---|
| NEXT 70 WDG herbicide at 21 g/ha (15 g a.e./ha) will control only: | |
| Lamb's-quarters* | Volunteer spring wheat |
| Stinkweed | (non-imazamox tolerant wheat) |
| Wild mustard | |
| NEXT 70 WDG herbicide at 29 g/ha (20 g a.e./ha) will control: | |
| Barnyard grass | Volunteer barley |
| Cleavers* | Volunteer canary seed |
| Cow cockle | Volunteer canola (non- imazamox tolerant canola only) |
| Green foxtail | Volunteer durum wheat |
| Green smartweed | Volunteer spring wheat (non-imazamox tolerant wheat) |
| Japanese brome grass* | Volunteer tame oats |
| Kochia* | Wild buckwheat* |
| Lamb's-quarters | Wild mustard |
| Persian dandelion | Wild oats |
| Redroot pigweed | Yellow foxtail |
| Round-leaved mallow* | |
| Russian thistle | |
| Shepherd's purse | |
| Stinkweed | |

* = suppression only

Apply in 100 L/ha of water with MERGE® Adjuvant at a rate of 0.5% v/v (e.g. 5 L of MERGE Adjuvant per 1000 L spray solution).

Imazamox tolerant Canola

Application should be made from the 2 to 6 true leaf stage of the canola crop and after the weeds have emerged. To control broadleaf weeds, apply NEXT 70 WDG herbicide from the cotyledon to the 4-leaf stage. To control grasses, apply NEXT 70 WDG herbicide from the 1 to 4-true leaf stage until early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

Imazamox tolerant Canola Quality *Brassica juncea*

Application should be from the 2 to 6 true leaf stage of the crop and after the weeds have emerged. To control broadleaf weeds, apply NEXT 70 WDG herbicide from the cotyledon to the 4-leaf stage. To control grasses, apply NEXT 70 WDG herbicide from the 1 to 4 true leaf stage until early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

Imazamox tolerant Sunflowers

Application should be made from the 2 to 8 leaf stage of the sunflower crop and after the weeds have emerged. To control broadleaf weeds, apply NEXT 70 WDG herbicide from the cotyledon up to the 4-leaf stage. To control grasses, apply NEXT 70 WDG herbicide from the 1 to 4-true

leaf stage up until early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

Imazamox tolerant Lentils

Application should be made from the 2 to 6 leaf stage of the lentil crop and after the weeds have emerged. To control broadleaf weeds, apply NEXT 70 WDG herbicide from the cotyledon up to the 4-leaf stage. To control grasses, apply NEXT 70 WDG herbicide from the 1 to 4-true leaf stage up until early tillering.

FOR FIELD PEAS

Application should be made from the 1 to 6 true leaf stage and after weeds have emerged. Apply when broadleaf weeds are from the cotyledon to 4-leaf stage and when grassy weeds are at the 1 to 4 true leaf or early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.

TANK-MIX COMBINATION

When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions, and grazing limitations for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

NEXT 70 WDG Herbicide + Lontrel SG Soluble Granular Herbicide in imazamox tolerant Canola

NEXT 70 WDG herbicide may be tank-mixed with Lontrel SG soluble granular herbicide in imazamox tolerant Canola.

| Timing | Early post-emergence | | | | | | | |
|----------------------------|---|--|----------------|---------------------------|----------------|----------------|--------------------|-----------------------|
| Rate | 29 g/ha NEXT 70 WDG Herbicide + 100 g/ha Lontrel SG Granular Herbicide | | | | | | | |
| Water Volume | 100 L/ha | | | | | | | |
| Surfactant/Adjuvant | Add MERGE Adjuvant - 0.5% v/v | | | | | | | |
| Weeds Controlled | <p>This tank-mix combination will provide control of weeds as listed on the NEXT 70 WDG Herbicide. Refer to Broadleaf weed and Grass control Section.</p> <p>Improved weed control with the tank mix vs. NEXT 70 WDG Herbicide alone.</p> <table border="1" data-bbox="631 1608 1373 1736"> <thead> <tr> <th>Control</th> <th>Top Growth Control</th> </tr> </thead> <tbody> <tr> <td>Wild buckwheat</td> <td>Canada thistle</td> </tr> <tr> <td>Annual sow thistle</td> <td>Perennial sow thistle</td> </tr> </tbody> </table> | | Control | Top Growth Control | Wild buckwheat | Canada thistle | Annual sow thistle | Perennial sow thistle |
| Control | Top Growth Control | | | | | | | |
| Wild buckwheat | Canada thistle | | | | | | | |
| Annual sow thistle | Perennial sow thistle | | | | | | | |

| | |
|---------------|--|
| Remark | <p>Application should be made from the 2 to 6 leaf stage of the canola crop and after the weeds have emerged.</p> <p>To control broadleaf weeds, apply NEXT 70 WDG herbicide from the cotyledon to the 4-leaf stage. To control grasses, apply NEXT 70 WDG herbicide from the 1 to 4 true leaf stage until early tillering. Initial transient crop yellowing may be observed after application but this is outgrown and should not affect yield.</p> <p>For tank mixing, see mixing instructions.</p> <p>Refer to Tank mix rotational crop restrictions for details.</p> |
|---------------|--|

Tank Mix Rotational Crop Restrictions

NEXT 70 WDG Herbicide + Lontrel SG Soluble Granular Herbicide in Imazamox Tolerant Canola

Fields previously treated with tank mix can be seeded to wheat, barley, oats and rye (not underseeded to forage legumes, clover or alfalfa), canola, corn, field peas, flax, forage grasses or can be summer-fallowed. Seed only those crops listed above in the year following treatment.

Do not seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application, delay seeding field peas an additional 12 months (total 22 months following application).

Contact your local Sharda CropChem Limited representative or retailer for more information before seeding field peas following drought conditions in the previous year.

Tank Mix Pre-harvest / Grazing Interval

Do not harvest the treated crop within 60 days of application.

BROADLEAF WEED and GRASS CONTROL (Eastern Canada Only - Soybeans)

Application must be made after the weeds have emerged. To control broadleaf weeds, apply NEXT 70 WDG Herbicide from the cotyledon to 4 leaf stage. To control grasses, apply NEXT 70 WDG Herbicide to green and yellow foxtail when in the 1-4 true leaf stage and to barnyard grass and fall panicum when in the 3-6 leaf stage.

NEXT 70 WDG Herbicide at 36 g product/ha will control:

- Common ragweed*
- Lady's thumb
- Lamb's-quarters*
- Redroot pigweed
- Velvetleaf
- Wild mustard
- Cocklebur*
- Eastern black nightshade
- Green foxtail

Yellow foxtail
Barnyard grass
Fall panicum

*Suppression only

Apply in 200 L/ha of water with a NONIONIC SURFACTANT (AGRAL[®], AGSURF[®] or SURF[®] 92) at the rate of 0.25% percent v/v and a fertilizer solution, 28-0-0 or 32-0-0, at a rate of 2 L/ha.

MIXING INSTRUCTIONS

1. Use 100 L/ha of water.
2. Use a 50-mesh (or coarser) filter screen.
3. Fill the spray tank with three-quarters of the required amount of clean water, start agitation and continue agitation throughout the entire mixing and spraying procedure.
4. Add the required amount of NEXT 70 WDG herbicide soluble bag(s) directly into the sprayer through the tank opening.
5. For NEXT 70 WDG Herbicide + Lontrel SG soluble granular herbicide tank mix, add recommended Lontrel SG soluble granular herbicide.
6. Agitate for at least ten minutes to dissolve the herbicide. After the herbicide is dissolved, continue agitation.
7. If required, add the required amount of MERGE Adjuvant. If excess foaming occurs, a silicone anti-foaming agent may be added (e.g. HALT[®]).
8. Complete filling the tank to the desired level with water. If agitation is stopped for more than 5 minutes, re-suspend spray solution by full agitation prior to commencing spraying again.
9. Between loads of NEXT 70 WDG Herbicide and Lontrel SG soluble granular herbicide, check in-line and nozzle screens and rinse clean if necessary.
10. Upon completion of spraying NEXT 70 WDG herbicide or the tank mix, thoroughly flush tank, boom, hoses and in-line and nozzle screens with clean water to avoid possible injury to other crops.

FOLLOW CROPPING

Winter wheat can be planted 3 months after treatment (3 MAT) as a rotational crop.

Initial crop injury to non-imazamox tolerant canola may be observed. Avoid spray overlap as yield reduction may result. The following crops may be grown safely the year following an application:

Canary seed
Field corn
Field peas
Imazamox tolerant Canola
Canola
Lentils
Spring wheat
Durum wheat
Spring barley
Sunflower
Tame oats

Flax
Chickpeas
Imazamox tolerant Sunflowers

The following crop may be grown safely two years following an application:

Mustard (condiment type only)

There are insufficient data for other follow crops. Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop other than those listed above.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, NEXT 70 WDG herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to NEXT 70 WDG herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of NEXT 70 WDG herbicide or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Sharda CropChem Limited at 1-844-810-5720 or 1-416-840-5639.

All products listed are registered trademarks of their respective companies.

Sleeve

| | | |
|-------|---|-----------|
| GROUP | 2 | HERBICIDE |
|-------|---|-----------|

NEXT 70 WDG HERBICIDE

Water Dispersible Granular Herbicide
in water-soluble bags

COMMERCIAL - AGRICULTURAL

ACTIVE INGREDIENT: Imazamox 70% a.e.

REGISTRATION NO. 33620 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 50 g to 20 kg (117.5 g water-soluble bags)

WARNING - EYE IRRITANT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

DO NOT SELL INDIVIDUAL WATER-SOLUBLE BAGS SEPARATELY

Sharda Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
Sharda Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, Canada
M9R1V1
1-844-810-5720
1-416-840-5639

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT APPLY using aerial application equipment.

Skin irritant.

WARNING - EYE IRRITANT

May cause eye damage.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, goggles or face shield, socks and shoes during mixing, loading, clean-up and repair. Goggle or faceshield are not required during application and gloves are not required during application within a closed cab.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

STORAGE

Store above 5°C.

Store in a cool, dry place away from children, animals, food, feed or fertilizers.

Keep unused water-soluble bags in resealed original container.

Keep packages dry at all times.

Keep packages out of rain.

Keep from freezing.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Pouch label

NEXT 70 WDG HERBICIDE

70% Water Dispersible Granular Herbicide

COMMERCIAL - AGRICULTURAL

NET CONTENTS: 117.5 g

ACTIVE INGREDIENT: Imazamox ... 70% active (a.e.)

WARNING - EYE IRRITANT

KEEP OUT OF REACH OF CHILDREN