

HERBICIDE

RUSH M™

This post-emergent herbicide gives you superior control of tough broadleaf weeds like chickweed, cleavers and wild buckwheat in spring wheat, durum wheat and barley.



ACTIVE INGREDIENT

Fluroxypyr and MCPA ester

CHEMISTRY GROUP

Group 4

APPLICATION RATES AND PACKAGING

- 240 ml/ac fluroxypyr + 380 ml/ac MCPA ester 600 or 20 ac/case
- 1 x 4.8 L fluroxypyr jug + 1 x 7.5 L MCPA ester 600
- 76.8 L fluroxypyr and 120 L MCPA ester treats 320 acres

REGISTERED CROPS:

- Barley
- Wheat (spring, durum)

WEEDS CONTROLLED:

2–4 leaf stage unless otherwise noted:

- Burdock
- Cleavers (1–4 whorls)
- Cocklebur
- Common ragweed
- Hempnettle (2–6 leaf)
- Kochia (including ALS-resistant biotypes)
- Lamb's quarters
- Mustards (except dog and tansy)
- Prickly lettuce
- Redroot pigweed
- Shepherd's purse
- Stinkweed
- Sunflower (annual)
- Vetch
- Volunteer canola
- Volunteer flax (½–5 inches)
- Wild mustard
- Wild radish

WEEDS SUPPRESSED:

- Smartweed (green)
- Stork's bill (1–8 leaf)
- Wild buckwheat (1–8 leaf)

HOW IT WORKS:

Systemic Group 4 herbicides disrupt normal plant growth regulation, resulting in death of susceptible plants.

CROP STAGING:

3 leaf up to the full emergence of the flag leaf.

WATER VOLUME:

40 L/ac

RAINFASTNESS:

None specified.

RUSH M™

SUPPORTED TANK MIXES:

Wheat and barley:

- Assert®
- Fenoxaprop
- Clodinafop (Ladder® 240 EC, Ladder® All-In) (wheat only)
- Tralkoxydim (Bison® 400 L)

MIXING INSTRUCTIONS:

1. Fill the spray tank ½ full with water and start agitation.
2. Add the required amount of Rush M™ and continue agitation.
3. Add the required amount of MCPA and continue agitation.
4. Fill tank with sufficient water to spray 40 L/ac of mixture.

CROP ROTATIONS:

Fields can be summerfallowed, or the following crops can be grown 1 year after application:

- | | |
|----------------|-----------|
| · Barley | · Mustard |
| · Canola | · Oats |
| · Flax | · Peas |
| · Forage grass | · Rye |
| · Lentils | · Wheat |

PRE-HARVEST INTERVAL:

60 days

GRAZING RESTRICTIONS:

- Do not cut hay within 7 days of application.
- Do not permit any grazing within 7 days of application.

STORAGE:

- Store in original container in dry, heated storage.
- If frozen, bring to room temperature and agitate before use.

QUICK TIPS:

The optimal temperature range is 12–24 C. Reduced activity will occur when temperatures are below 8 C or above 27 C. Frost 3 days before or after application may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions.