



# QUICK SHEET FOR TOMATO

<b>RATE</b>	35 - 70 g/ha (14 - 28 g/acre) + Non-ionic Surfactant	<b>DIRECT SEEDED</b>	<b>POST-emerge</b>	<ul style="list-style-type: none"> <li>• Apply between 4-leaf stage through to 1st bloom</li> <li>• Following bloom use only a directed spray or crop shields</li> </ul>
			<b>POST-emerge followed by POST-emerge</b>	<ul style="list-style-type: none"> <li>• Spot treatment only</li> <li>• Maximum of 70 g SANDEA per treated area</li> <li>• Allow 21 days between applications</li> </ul>

<b>RATE</b>	35 - 70 g/ha (14 - 28 g/acre) + Non-ionic Surfactant	<b>TRANSPLANTED</b>	<b>Bare Ground</b>	<p><b>Pre-transplant</b></p> <ul style="list-style-type: none"> <li>• Allow 7 days between application and transplanting</li> </ul> <p><b>Post-transplant</b></p> <ul style="list-style-type: none"> <li>• Apply before 1st bloom</li> <li>• Allow 14 days between application and transplanting</li> <li>• Following bloom use only a directed spray or crop shields</li> <li>• Apply to transplants that are established and actively growing</li> </ul>
			<b>Plastic Mulch</b>	<p><b>Pre-transplant</b></p> <ul style="list-style-type: none"> <li>• Apply following bed shaping and just before installation of plastic</li> <li>• Allow 7 days between application and transplanting</li> </ul>
			<b>PRE-emerge followed by POST-emerge</b>	<ul style="list-style-type: none"> <li>• Spot treatment only</li> <li>• Maximum of 47 g SANDEA per treated area</li> </ul>
			<b>POST-emerge followed by POST-emerge</b>	<ul style="list-style-type: none"> <li>• Spot treatment only</li> <li>• Maximum of 70 g SANDEA per treated area</li> <li>• Allow 21 days between applications</li> </ul>

<b>WEEDS</b>	<ul style="list-style-type: none"> <li>• SANDEA will provide control or suppression of over 40 weeds but is especially strong on yellow nutsedge, common cocklebur, galinsoga, wild mustard, redroot &amp; smooth pigweed, wild radish, common ragweed, sunflower and velvetleaf.</li> <li>• Weed spectrum may be further enhanced with the use of an appropriate tank mix partner</li> <li>• Control of weeds is optimal if weeds are treated while small and actively growing.</li> <li>• Rate is dependent on size of weed at time of application. See weeds table in label for details.</li> <li>• <b>May not control ALS (Group 2) resistant weeds</b></li> <li>• Within 48 hours of application of SANDEA growth of susceptible weeds is inhibited and susceptible weeds are no longer competitive with crop.</li> </ul>
--------------	--



# QUICK SHEET FOR **TOMATO**

## NOTES

- Pre-harvest Interval: 30 days
- Water Volume: 190 L/ha (20 gallons/acre)
- Use lower rate on lighter textured soils with low organic matter
- Apply **at least** 7 days before transplanting or 14 days following seeding/transplanting
- Treated soil in transplant hole may result in crop injury
- **DO NOT** spray on top of plastic mulch as spray solution may concentrate in hole and cause crop injury
- Maximum of 2 applications per season Not all varieties have been tested.
- End-user must consider the potential for a delayed harvest **BEFORE** using SANDEA as:
  - Applications of SANDEA may cause temporary stunting and/or delayed maturity which may result in delayed harvest
  - Under adverse growing conditions (dry or excessive moisture, cool weather, etc) maturity of the crop may be delayed which can influence harvest date, yield and quality
- To avoid injury to desirable crops, **CLEAN ALL MIXING AND SPRAY EQUIPMENT** before and immediately following applications of SANDEA.

## BUFFER ZONES\*

Fresh Water		Terrestrial Habitat
Less than 1 m	Greater than 1 m	
15 m	5 m	30 m

\* The spray drift buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site (<http://www.hc-sc.gc.ca/cps-spc/pest/agri-commerce/drift-derive/calculator-calculatrice-eng.php>). Always read and follow all label instructions. PMRA Registration# 31209 **Canyon Group L.L.C.** Sandeia™ is a trademark of Gowan Company, L.L.C.

Gowan Company recommends the following re-cropping intervals for crop safety. Planting prior to the intervals shown below may result in crop injury when using SANDEA Herbicide. Rotation intervals below may need to be extended if drought or cool conditions prevail. **Refer to individual product labels to determine rotational crop restrictions when tank mixtures are used. Refer to SANDEA label for additional information.**

## CROP ROTATION INFORMATION

<b>0 Months</b> (immediate plantback)	• Beans ( <i>dry</i> )
<b>1 Month</b>	• Corn, field normal and all herbicide tolerant varieties
<b>2 Months</b>	• Cereals, spring ( <i>barley, oats, wheat</i> ) • Cereals, winter ( <i>barley, wheat, rye</i> ) • Corn, seed • Forage grasses • Proso millet • Sorghum
<b>3 Months</b>	• Corn, sweet and pop
<b>6 Months</b>	• Peanuts
<b>8 Months</b>	• Tomato
<b>9 Months</b>	• Beans ( <i>snap</i> ) • Cucumbers • Forage Legumes ( <i>alfalfa, clovers</i> ) • Melons • Peas ( <i>succulent, field</i> ) • Potatoes • Pumpkins • Soybean • Squash
<b>10 Months</b>	• Peppers
<b>12 Months</b>	• Eggplant • Radish
<b>15 Months</b>	• Cabbage • Canola • Carrot • Mint
<b>18 Months</b>	• Broccoli • Cauliflower • Collards • Lettuce • Onions and Leeks • Sunflowers
<b>24 Months</b>	• Spinach
<b>36 Months</b>	• Strawberries • Sugarbeets • Table ( <i>garden</i> ) beets

For more information, contact your local Gowan Agro Canada representative: **Patti Turner 289-922-8174**