



Bromotril® II

Bromoxynil

Herbicide – Pre-Seed Burndown

When tank-mixed with glyphosate, Bromotril II provides outstanding burndown control of tough weeds like kochia, buckwheat, cow cockle and Russian thistle

Active ingredient: Bromoxynil

Chemistry group: Group 6

How it works: Bromotril II is a contact herbicide. Leaves will yellow in 2 to 4 days with complete control in 7 to 14 days. Bromotril II controls Group 2 and Group 9 (glyphosate) resistant biotypes.

Registered crops: Tank-mixed with glyphosate, this pre-seed burndown should only be used for minimum or zero-tillage cropping systems in wheat, barley and oat

Crop staging:

- Application can be made just prior to or immediately after seeding
- Under no circumstances should application be made after crop emergence

Weeds controlled:

Seedling up to 4 leaf stage:

- | | | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none">• American nightshade• Bluebur• Cocklebur• Common ragweed• Cow cockle¹ | <ul style="list-style-type: none">• Green smartweed• Kochia²• Lady's-thumb• Pale smartweed• Pigweed¹ | <ul style="list-style-type: none">• Russian thistle²• Stinkweed¹• Velvetleaf³• Wild mustard¹ |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

Seedling up to 8 leaf stage:

- | | |
|------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none">• Common buckwheat• Common groundsel• Lamb's-quarter | <ul style="list-style-type: none">• Tartary buckwheat• Wild buckwheat |
|------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------|

Grassy weeds up to 15 cm:

- Green foxtail
- Volunteer cereals
- Wild Oats

¹ In normal conditions, will be controlled up to 4 leaf stage. Plants beyond this stage are unlikely to be controlled; the higher rate generally gives better results

² Spray before plants are 5 cm high. Including Group 2 and Group 9 resistant biotypes

³ Spray before plants are 8 cm high





Application rates and packaging:

- Bromotril II at 0.49 L/acre (40 ac/case) tank-mixed with glyphosate at a rate of 175g a.i. per acre
- 2 x 9.7L jugs/case

Water volume:

- Ground: 20 – 40 L/ac

Rainfastness: 30 minutes

Registered and supported tank mixes:

- Must be tank-mixed with glyphosate at a rate of 175g a.i. per acre of glyphosate

Mixing instructions:

1. Fill spray tank half full with clean water
2. Add required amount of Bromotril II. Begin agitation
3. Add the recommended amount of glyphosate to the spray tank, agitate
4. Add surfactant
5. Complete filling spray tank with water, continue agitation

Crop rotations: Only apply prior to wheat, barley and oat

Storage: Avoid freezing

Quick tips:

- For best results, spray when weeds are in the seedling stage

