



Outshine™

Florasulam, Fluroxypyr & MCPA Ester

Herbicide

Post-emergent broadleaf herbicide for control of hard-to-kill annual weeds including Group 2 resistant cleavers, hemp-nettle and kochia in spring wheat, durum wheat and spring barley

Active ingredients: Florasulam, fluroxypyr and MCPA ester

Chemistry group: Group 2 (florasulam) and Group 4 (fluroxypyr and MCPA ester)

How it works: Outshine, applied early and thoroughly to the main flush of actively growing broadleaved weeds, quickly causes plants to stop growing, even if typical symptoms of dying weeds are not noticeable for 1 to 2 weeks after application.

Registered crops:

- Barley
- Wheat (spring and durum)

Weeds controlled:

- | | | |
|---|--------------------|----------------------|
| • Burdock | • Prickly lettuce | • Sunflower (annual) |
| • Chickweed (common) | • Ragweed | • Vetch |
| • Cleavers | • Redroot pigweed | • Volunteer canola |
| • Cocklebur | • Russian pigweed | • Volunteer flax |
| • Hemp-nettle | • Shepherd's-purse | • Wild buckwheat |
| • Kochia (including ALS resistant biotypes) | • Smartweed | • Wild mustard |
| • Lamb's-quarters | • Stinkweed | • Wild radish |
| • Plantain | • Stork's-bill | |

Crop staging: 2 leaf expanded to 6 leaf stage

Application rates and packaging:

- 40 ac/case
- 2 x 8L florasulam + fluroxypyr and 1 x 9.33L MCPA

Water volume: Ground: minimum 40 L/ac

Rainfastness: 2 hours


Registered and supported tank mixes:

Barley

- Assert®
- Axial®

Wheat (spring and durum)

- Assert®
- Axial® (spring wheat only)
- Everest®





Herbicide

Outshine™

Florasulam, Fluroxypyr & MCPA Ester



Mixing instructions:

1. Fill spray tank half full with clean water and begin agitation
2. Add the required amount of Outshine
3. Continue filling the tank with sufficient water to spray 40L of spray mixture per acre
4. Use caution near susceptible crops or desirable plants
5. Product has the potential to leach: avoid excessive irrigation

Crop rotations:

- Can be seeded the following year to barley, canola, oats, peas or wheat, or fields can be summerfallowed
- Do not use in successive years at the same site

Pre-harvest interval: Do not harvest treated crop within 60 days of application

Grazing restriction: Do not graze livestock within 7 days of application

Storage: Store in original containers in secure, dry, heated storage. If product is frozen, bring to room temperature and agitate before use

Quick tips:

- Two unique modes of action and three active ingredients provide resistance management and high performance



Simply. Grow. Together.

ADAMA

Always read and follow label directions
Toll-free: 1.855.264.6262
WEBSITE: adama.com/canada

™ Outshine is a trademark of ADAMA Agricultural Solutions Canada Ltd.
All others are trademarks of their respective companies. 176-1/0215