



Rush M™

Fluroxypyr & MCPA Ester



Herbicide

Post-emergent herbicide for superior control of tough broadleaf weeds like chickweed, cleavers and wild buckwheat, in wheat (spring and durum) and barley

Active ingredients: Fluroxypyr and MCPA ester

Chemistry group: Group 4

How it works: Systemic Group 4 herbicides disrupt normal plant growth regulation resulting in death of susceptible plants.

Registered crops:

- Barley
- Wheat (spring and durum)

Weeds controlled:


2 to 4 leaf stage unless otherwise noted:

- | | | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none">• Burdock• Cleavers (1-4 whorls)• Cocklebur• Flixweed• Hemp-nettle (2-6 leaf)• Kochia• Lamb's-quarters | <ul style="list-style-type: none">• Mustards (except dog and tansy)• Prickly lettuce• Ragweed (common)• Redroot pigweed• Shepherd's-purse• Stinkweed | <ul style="list-style-type: none">• Sunflower (annual)• Vetch• Volunteer canola• Volunteer flax (1-12 cm)• Wild mustard• Wild radish |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

Weeds suppressed:

- Smartweed (green)
- Stork's-bill (1-8 leaf)
- Wild buckwheat (1-8 leaf)

Crop staging: 3 leaf up to the full emergence of the flag leaf





Application rates and packaging:

- 240 mL/ac fluroxypyr + 380 mL/ac MCPA ester 600, or 20 acres/case
- 1 x 4.8L fluroxypyr jug + 1 x 7.5L MCPA ester 600/case

Water volume: 40 L/ac

Rainfastness: None specified

Registered and supported tank mixes:

Wheat and barley:

- Assert®
- Fenoxaprop (Bengal®)
- Clodinaflop (Ladder®)
- Tralkoxydim (Bison®)

Mixing instructions:

1. Fill the sprayer tank half full with water and start agitation running
2. Add the required amount of Rush M and continue agitation
3. Add the required amount of MCPA and continue agitation
4. Fill tank with sufficient water to spray 40L of mixture per acre

Crop rotations:

The following crops can be grown one year after application, or fields can be summerfallowed:

- Barley
- Canola
- Flax
- Forage grass
- Lentil
- Mustard
- Oat
- Pea
- Rye
- Wheat

Pre-harvest interval: 60 days

Grazing restriction:

- Do cut hay within 7 days of application
- Do **NOT** permit any grazing within 7 days of application

Storage:

- Store in original container in dry, heated storage
- If frozen, bring to room temperature and agitate before use

Quick tips:

- Optimum temperature range 12°C to 24°C
- Reduced activity will occur when temperatures are below 8°C or above 27°C
- Frost 3 days before or after application may reduce weed control and crop tolerance
- Weed control may be reduced during stress conditions

ADAMA

Always read and follow label directions
Toll-free: 1.855.264.6262
WEBSITE: adama.com/canada



Simply. Grow. Together.

® Bengal, Bison and Ladder are registered trademarks and ™ Rush M is a trademark of ADAMA Agricultural Solutions Canada Ltd. All others are trademarks of their respective companies. 176-1/0115