




Topnotch™

Azoxystrobin & Propiconazole



Fungicide

Unique azoxystrobin & propiconazole formulation to provide broad-spectrum protection against cereal leaf diseases

Active ingredient: Azoxystrobin and propiconazole

Chemistry group: Group 3 (propiconazole) and Group 11 (azoxystrobin)

How it works: Used as both a curative and preventative fungicide, Topnotch has broad spectrum, systemic and contact activity.

Registered crops:

- Barley
- Oats
- Rye
- Triticale
- Wheat

Key diseases controlled:

- Barley leaf rust
- Net and spot blotches
- Scald
- Septoria spot
- Stripe rust
- Tan spot
- Wheat leaf rust

Application timing and crop staging:

Crop	Diseases	Application Timing
Barley	Barley net blotch Barley scald Septoria leaf spot Stripe rust Barley leaf rust Tan spot	Apply once between stem elongation and half-head emergence (G.S. 29-55)
Oats	Barley net blotch Crown rust Septoria leaf spot	Apply once between stem elongation and half-head emergence (G.S. 29-55)
Rye	Septoria leaf spot Barley scald Tan spot	Apply once between stem elongation and half-head emergence (G.S. 29-55)
Triticale	Septoria leaf spot Tan spot	Apply once between stem elongation and half-head emergence (G.S. 29-55)
Wheat	Septoria leaf spot Tan spot Stripe rust Wheat leaf rust	Apply once between stem elongation and half-head emergence (G.S. 29-55)





Fungicide

Topnotch™

Azoxystrobin & Propiconazole



Application rates and packaging:

- Topnotch at 212 mL/ac or 80 ac/case
- 2 x 8.6L jugs/case

Water volume:

- Ground: minimum 40 L/ac
- Aerial: 18 L/ac

Rainfastness: None specified

Registered and supported tank mixes:

Herbicides

- Axial®
- Broadband®
- Clodinafop-propargyl (Ladder®, Horizon®)
- Sierra®
- Touchdown Total®
- Traxion™

Insecticides

- Lambda-cyhalothrin (Silencer®, Matador®)

Fungicides

- Quadris®

Mixing instructions:

1. Fill spray tank 1/2 to 2/3 full with water
2. With agitator running, add required amount of Topnotch and continue agitating while adding remainder of the water
3. Begin application after Topnotch is completely dispersed into the mix water, and maintain agitation during spraying operation

Crop rotations: Do not plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of application

Pre-harvest interval: 45 days

Grazing restriction: None known

Storage: Do not store below 0°C

Quick tips:

- Good spray coverage and canopy penetration are important to achieve the best results



Simply. Grow. Together.

® Ladder and Silencer are registered trademarks and ™ Topnotch is a trademark of ADAMA Agricultural Solutions Canada Ltd. All others are trademarks of their respective companies. 176-1/0215

ADAMA

Always read and follow label directions
Toll-free: 1.855.264.6262
WEBSITE: adama.com/canada