

Sword®

HERBICIDE

For systemic control of many broadleaf weeds in: barley, canary seed, oats, summerfallow, wheat, and seedling and established grasses grown for forage.

WEEDS CONTROLLED:

Buckwheat (wild, tartary, volunteer cultivated)	Mustard (wild, ball, tall, wormseed, yellow)
Cleavers*	Night flowering catchfly
Common ragweed	Pigweed (prostrate, redroot)
Corn spurry	Russian thistle, ***
Cow cockle	Shepherd's purse
Flixweed	Annual sow-thistle,
Green smartweed	Stinkweed
Hemp-nettle**	Volunteer rapeseed
Knotweed	Volunteer sunflower
Kochia	Also chickweed, wild oats, and millet by tank mixes.
Lady's-thumb	
Lamb's-quarters	

TOP GROWTH CONTROL ONLY:

Canada thistle	Hedge bindweed
Field bindweed	Perennial sow-thistle

* Cleavers: spray in 1-2 whorl stage.

** Hemp-nettle: spray before second pair of true leaves.

*** Russian thistle: spray when less than 5 cm in height.

CROPS: Barley, canary seed, durum wheat, oats, summerfallow, wheat, winter wheat, and seedling and established grasses grown for forage. (Not underseeded to legumes.)

RATE: Apply 1.0-1.5 L/ha (.4-.6 L/acre) (one 10 L jug treats 25 to 16.7 acres).

WATER VOLUME: Use 50-100 L/ha (5-20 gal/acre) by ground and minimum of 30 L/ha (3 gal/acre) by air.

TANK MIXES: Horizon®, Everest, (wheat only). Nortrace plant nutrients. (Conduct a jar test with Nortrace plant nutrients before tank mixing.) For other tank mixes see label.

TIMING:

Barley: 2 to 4 leaf.

Wheat: (spring, durum), oats, canary seed, 2 to 5 leaf.

Winter Wheat: apply in the spring before the crop is more than 30 cm high (top leaf extended).

WINTER ANNUAL WEEDS: Should be sprayed as soon as new growth starts.

CAUTION: Best results are obtained when weeds are in the 2-3 leaf stage and actively growing. Less than satisfactory control may result if weeds are not actively growing.



