

BASE

LAGON® 480 E

INSECTICIDE

COMMERCIAL

This product is not to be used in and around homes or other residential areas such as parks, school grounds, playing fields. It is not for use by homeowners or other uncertified users.



WARNING

POISON

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Dimethoate 480 g/L

REGISTRATION NO. 9382 PEST CONTROL PRODUCTS ACT

IN CASE OF EMERGENCY DUE TO A MAJOR SPILL, FIRE OR POISONING
INVOLVING THIS PRODUCT CALL DAY OR NIGHT, 1-800-561-8273

LOVELAND PRODUCTS CANADA INC.
789 Donnybrook Drive
Dorchester, ON
N0L 1G5
1-800-328-4678

NET CONTENTS: 1 L – 115 L

2020-0144
2021-02-11

GROUP	1B	INSECTICIDE
-------	-----------	-------------

FRONT

LAGON® 480 E

INSECTICIDE

COMMERCIAL

This product is not to be used in and around homes or other residential areas such as parks, school grounds, playing fields. It is not for use by homeowners or other uncertified users.



WARNING

POISON

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Dimethoate 480 g/L

REGISTRATION NO. 9382 PEST CONTROL PRODUCTS ACT

IN CASE OF EMERGENCY DUE TO A MAJOR SPILL, FIRE OR POISONING
INVOLVING THIS PRODUCT CALL DAY OR NIGHT, 1-800-561-8273

LOVELAND PRODUCTS CANADA INC.
789 Donnybrook Drive
Dorchester, ON
N0L 1G5
1-800-328-4678

NET CONTENTS: 1 L – 115 L

1/16

2020-0144
2021-02-11

PRECAUTIONS

**HARMFUL OR FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN
KEEP OUT OF REACH OF CHILDREN**

Keep the following personal protective equipment immediately available for use in case of emergency (for example, a broken package, spill, or equipment breakdown): chemical-resistant coveralls, chemical-resistant gloves, chemical-resistant head gear and a respirator.

Not for use in greenhouses.

This product is not to be used around homes or other residential areas such as parks, school grounds and/or playing fields. It is not for use by homeowners or other uncertified users.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Avoid breathing vapour or spray mist. Use only with adequate ventilation. Do not use indoors. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling and before eating and smoking. Avoid contamination of feed and food stuffs. Keep away from heat, sparks and open flame.

Do not use in milk processing or storage rooms. Remove livestock and poultry from buildings when spraying. Do not contaminate feed troughs, litter or drinking fountains. Do not use indoors. Do not mix with whitewash or apply within 2 weeks of whitewashing. Keep out of sun.

When spraying honey-producing crops (alfalfa, red clover, sweet clover and alsike) spray at least 5 days before bloom appears and do not introduce hives until full bloom. Hives should be removed from vicinity before making spray applications. If spraying must be done during the blooming period, restrict application to the period after dark or very early morning in order to reduce mortality to bees that may be visiting the blooms.

This product is toxic to birds and wildlife. To reduce injury to bees, restrict application to the period after dark when bees are inside the hives or in the early morning before the bees are foraging in the fields. Do not apply to such crops as alfalfa when in full bloom.

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.

Ship and store between 4° and 30° C.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact LOVELAND PRODUCTS CANADA INC.

Engineering Controls and Personal Protective Equipment:

A. Mixing/loading liquids (for all uses except forestry):

Wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves, goggles or a face shield and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides or a NIOSH approved canister approved for pesticides.

B. Mixing/loading liquids (forestry: hemlock, spruce and balsam fir):

Use a closed system for aerial mixing/loading for forestry use (hemlock, spruce and balsam fir). Wear cotton coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves, goggles or a face shield.

C. Mixing/loading liquids (forestry: Douglas-fir (seed tree), Sitka spruce (seed tree), spruce (seed tree and woodland)):

During mixing, loading, clean-up and repair, wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves, goggles or a face shield and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides or a NIOSH approved canister approved for pesticides.

D. Applying by air:

Wear cotton coveralls over long pants, and a long-sleeved shirt, shoes, plus socks.

E. Applying by groundboom:

During groundboom application use a closed cab that provides both a physical barrier and respiratory protection (for example, dust/mist filtering and/or vapour/gas purification system). The closed cab must have a chemical resistant barrier that totally surrounds the occupant and prevents contact with pesticides outside the cab. Wear cotton coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks. Chemical resistant gloves are not required to be worn while driving closed cab equipment but are required for clean-up and repair activities.

2020-0144
2021-02-11

F. Applying by airblast:

During airblast application use a closed cab that provides both a physical barrier and respiratory protection (for example, dust/mist filtering and/or vapour/gas purification system). The closed cab must have a chemical resistant barrier that totally surrounds the occupant and prevents contact with pesticides outside the cab. Wear cotton coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks. Chemical resistant gloves are not required to be worn while driving closed cab equipment but are required for clean-up and repair activities.

If a closed cab is not feasible, wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves and chemical-resistant headgear. Chemical resistant headgear includes sou'westers, or large brimmed, water-proof hats, and hoods with sufficient neck protection. Avoid touching face or other unprotected parts of the body during application.

G. Applying by handheld equipment:

Wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides or a NIOSH approved canister approved for pesticides. Limit the amount of active ingredient handled per day to 2.8 kg per person (for example, approximately 2950 L at a rate of 0.96 g a.i./L) when applying by handheld equipment.

H. Applying by manually-pressurized handwand or backpack (Douglas-fir (seed tree), Sitka spruce (seed tree), spruce (seed tree and woodland)):

During manually-pressurized handwand, or backpack application, wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH approved canister approved for pesticides.

DO NOT apply using mechanically-pressurized handgun equipment. Only for use with manually-pressurized hand wands or backpack sprayers.

I. Applying by right-of-way sprayer:

During mixing, loading, application, clean-up and repair, wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical-resistant gloves, goggles or a face shield.

J. Applying by soil drench, soil injection or chemigation:

Wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides or a NIOSH approved canister approved for pesticides.

ENVIRONMENTAL PRECAUTIONS:

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment. **TOXIC** to bees. Bees may be exposed through direct spray, spray drift, and residues on leaves, pollen and nectar in flowering crops and weeds. Minimize spray drift to reduce harmful effects on bees in habitats close to the application site. Avoid applications when bees are foraging in the treatment area in groundcover containing blooming weeds. To further minimize exposure to pollinators, refer to the complete guidance "Protecting Pollinators during Pesticide Spraying – Best Management Practices" on the Health Canada website (www.healthcanada.gc.ca/pollinators). Follow crop specific directions for application timing.

For applications on crops that are highly attractive to pollinators (alfalfa, clovers, canola, safflower, blueberries, cherries, peaches, pears, asparagus, and outdoor ornamentals excluding coniferous evergreens), or when using managed bees for pollination services:

DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

For applications on all other crops:

Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging.

TOXIC to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.

TOXIC to birds, mammals and aquatic organisms. Observe spray buffer zones specified under **DIRECTIONS FOR USE**.

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (for example, soils that are compacted or fine textured such as clay).

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

DECONTAMINATION

If accidental spillage of LAGON 480 E Insecticide should occur, scrub contaminated area immediately with a strong laundry soap solution or use household lye. Detergents are not satisfactory for this purpose. Repeated scrubblings are necessary on plain wood surfaces. Avoid spillage on all types of floor tile or linoleum.

2020-0144
2021-02-11

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Dimethoate is an organophosphate that is a cholinesterase inhibitor. Typical symptoms of overexposure to cholinesterase inhibitors include headache, nausea, dizziness, sweating, salivation, and runny nose and eyes. This may progress to muscle twitching, weakness, tremor, incoordination, vomiting, abdominal cramps and diarrhea in more serious poisonings. A life-threatening poisoning is signified by loss of consciousness, incontinence, convulsions and respiratory depression with a secondary cardiovascular component. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate degree of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as pralidoxime chloride, may be therapeutic if used early; however, use only in conjunction with atropine. In cases of severe acute poisoning, use antidotes immediately after establishing an open airway and respiration. With oral exposure, the decision of whether to induce vomiting or not should be made by an attending physician. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

STORAGE

Do not use, pour, spill, or store near heat, sparks, or open flame. Keep out of direct sunlight. Ship and store between 4°C and 30°C.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

2020-0144
2021-02-11

GROUP	1B	INSECTICIDE
-------	-----------	-------------

DATAPAK

LAGON® 480 E

INSECTICIDE

COMMERCIAL

This product is not to be used in and around homes or other residential areas such as parks, school grounds, playing fields. It is not for use by homeowners or other uncertified users.



WARNING

POISON

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Dimethoate 480 g/L

REGISTRATION NO. 9382 PEST CONTROL PRODUCTS ACT

IN CASE OF EMERGENCY DUE TO A MAJOR SPILL, FIRE OR POISONING
INVOLVING THIS PRODUCT CALL DAY OR NIGHT, 1-800-561-8273

LOVELAND PRODUCTS CANADA INC.
789 Donnybrook Drive
Dorchester, ON
N0L 1G5
1-800-328-4678

NET CONTENTS: 1 L – 115 L

1/16

2020-0144
2021-02-11

PRECAUTIONS
HARMFUL OR FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN
KEEP OUT OF REACH OF CHILDREN

Keep the following personal protective equipment immediately available for use in case of emergency (for example, a broken package, spill, or equipment breakdown): chemical-resistant coveralls, chemical-resistant gloves, chemical-resistant head gear and a respirator.

Not for use in greenhouses.

This product is not to be used around homes or other residential areas such as parks, school grounds and/or playing fields. It is not for use by homeowners or other uncertified users.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Avoid breathing vapour or spray mist. Use only with adequate ventilation. Do not use indoors. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling and before eating and smoking. Avoid contamination of feed and food stuffs. Keep away from heat, sparks and open flame.

Do not use in milk processing or storage rooms. Remove livestock and poultry from buildings when spraying. Do not contaminate feed troughs, litter or drinking fountains. Do not use indoors. Do not mix with whitewash or apply within 2 weeks of whitewashing. Keep out of sun.

When spraying honey-producing crops (alfalfa, red clover, sweet clover and alsike) spray at least 5 days before bloom appears and do not introduce hives until full bloom. Hives should be removed from vicinity before making spray applications. If spraying must be done during the blooming period, restrict application to the period after dark or very early morning in order to reduce mortality to bees that may be visiting the blooms.

This product is toxic to birds and wildlife. To reduce injury to bees, restrict application to the period after dark when bees are inside the hives or in the early morning before the bees are foraging in the fields. Do not apply to such crops as alfalfa when in full bloom.

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.

Ship and store between 4° and 30° C.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact LOVELAND PRODUCTS CANADA INC.

Engineering Controls and Personal Protective Equipment:

A. Mixing/loading liquids (for all uses except forestry):

Wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves, goggles or a face shield and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides or a NIOSH approved canister approved for pesticides.

B. Mixing/loading liquids (forestry: hemlock, spruce and balsam fir):

Use a closed system for aerial mixing/loading for forestry use (hemlock, spruce and balsam fir). Wear cotton coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves, goggles or a face shield.

C. Mixing/loading liquids (forestry: Douglas-fir (seed tree), Sitka spruce (seed tree), spruce (seed tree and woodland)):

During mixing, loading, clean-up and repair, wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves, goggles or a face shield and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides or a NIOSH approved canister approved for pesticides.

D. Applying by air:

Wear cotton coveralls over long pants, and a long-sleeved shirt, shoes, plus socks.

E. Applying by groundboom:

During groundboom application use a closed cab that provides both a physical barrier and respiratory protection (for example, dust/mist filtering and/or vapour/gas purification system). The closed cab must have a chemical resistant barrier that totally surrounds the occupant and prevents contact with pesticides outside the cab. Wear cotton coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks. Chemical resistant gloves are not required to be worn while driving closed cab equipment but are required for clean-up and repair activities.

F. Applying by airblast:

During airblast application use a closed cab that provides both a physical barrier and respiratory protection (for example, dust/mist filtering and/or vapour/gas purification system). The closed cab must have a chemical resistant barrier that totally surrounds the occupant and prevents contact with pesticides outside the cab. Wear cotton coveralls over long pants and a long-sleeved shirt,

2020-0144
2021-02-11

chemical resistant footwear plus socks. Chemical resistant gloves are not required to be worn while driving closed cab equipment but are required for clean-up and repair activities.

If a closed cab is not feasible, wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves and chemical-resistant headgear. Chemical resistant headgear includes sou'westers, or large brimmed, water-proof hats, and hoods with sufficient neck protection. Avoid touching face or other unprotected parts of the body during application.

G. Applying by handheld equipment:

Wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides or a NIOSH approved canister approved for pesticides. Limit the amount of active ingredient handled per day to 2.8 kg per person (for example, approximately 2950 L at a rate of 0.96 g a.i./L) when applying by handheld equipment.

H. Applying by manually-pressurized handwand or backpack (Douglas-fir (seed tree), Sitka spruce (seed tree), spruce (seed tree and woodland)):

During manually-pressurized handwand, or backpack application, wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH approved canister approved for pesticides.

DO NOT apply using mechanically-pressurized handgun equipment. Only for use with manually-pressurized hand wands or backpack sprayers.

I. Applying by right-of-way sprayer:

During mixing, loading, application, clean-up and repair, wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical-resistant gloves, goggles or a face shield.

J. Applying by soil drench, soil injection or chemigation:

Wear chemical resistant coveralls over long pants and a long-sleeved shirt, chemical resistant footwear plus socks, chemical resistant gloves and suitable respiratory protection. Suitable respiratory protection is defined as either a respirator with a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides or a NIOSH approved canister approved for pesticides.

ENVIRONMENTAL PRECAUTIONS:

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.

TOXIC to bees. Bees may be exposed through direct spray, spray drift, and residues on leaves, pollen and nectar in flowering crops and weeds. Minimize spray drift to reduce harmful effects on bees in habitats close to the application site. Avoid applications when bees are foraging in the treatment area in groundcover containing blooming weeds. To further minimize exposure to pollinators, refer to the complete guidance "Protecting Pollinators during Pesticide Spraying – Best Management Practices" on the Health Canada website (www.healthcanada.gc.ca/pollinators). Follow crop specific directions for application timing.

For applications on crops that are highly attractive to pollinators (alfalfa, clovers, canola, safflower, blueberries, cherries, peaches, pears, asparagus, and outdoor ornamentals excluding coniferous evergreens), or when using managed bees for pollination services:

DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

For applications on all other crops:

Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging.

TOXIC to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.

TOXIC to birds, mammals and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (for example, soils that are compacted or fine textured such as clay).

Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

DIRECTIONS FOR USE

LAGON 480E INSECTICIDE is a true systemic insecticide. A systemic insecticide is absorbed into the system of the plant upon application and, as with all systemic materials, may in specific plants cause reactions which are neither predictable nor common to all members of the species.

LAGON 480 E Insecticide is generally effective in controlling aphids, and other listed insects, scales and mites.

LAGON 480 E Insecticide may be applied by spray or soil drench. LAGON 480 E Insecticide is an emulsifiable concentrate. Use designated amounts in full volume of water.

2020-0144
2021-02-11

SPRAY BUFFER ZONES:

Spot treatments using handheld equipment DO NOT require a buffer zone. Use of soil drench or soil incorporation do not require a spray buffer zone.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:			
			Freshwater habitat of Depth:		Estuarine/Marine habitats of Depth:	
			Less than 1m	Greater than 1m	Less than 1m	Greater than 1m
Field Sprayer*	Cereals(barley, oats, rye, wheat), forage crops (clover, grasses, hay, grain), alfalfa (forage and seed production), clover (sweet, red, alsike), pastures and wastelands, safflower, soybeans, bok choi (pak choi), chicory, Chinese broccoli, cole crops (brussel sprouts), peppers, potatoes, tomatoes		2	1	1	1
	Flax, canary grass (seed production)		1	0	1	0
	Canola (rape), beans (pole, snap), beets, celery, eggplant, leafy vegetables (beet greens, kale, leaf lettuce, Swiss chard, turnip greens), Blueberry (low and highbush)		1	1	1	1
	Asparagus		3	1	2	1
	Cole crops (broccoli, cauliflower)		2	1	2	1
	Peas (field and/or canning)		1	0	1	0
	Strawberries		3	2	3	1
	Christmas Trees: Balsam fir, hemlock, pine (Mugho, red, Scots), spruce; Ornamental trees: Arbor vitae, cedar, juniper, oak, pine (Mugho, red, Scots), Taxus (yew), Lilac (<i>Syringa</i> spp.)		2	1	2	1
	Birch		1	0	1	0
	Boxwood, hemlock, holly (English, American)		1	1	1	1
	Douglas-fir (seed tree), spruce (seed tree and woodland)		10	5	10	5
	Larch		3	2	4	2
	Sitka spruce (seed tree)		15	5	10	5
	Day lilies, irises, euonymus		2	1	2	1
	Roses, azaleas, gardenias, camellias, gerberas, gladioli, poinsettias		1	1	1	1
	Airblast	Christmas trees Balsam fir, hemlock, pine (Mugho, red, Scots), spruce; Ornamental trees: Arbour vitae, cedar, juniper, oak, pine (Mugho, red, Scots), Taxus (yew)	Late growth stage	4	2	4
Birch			1	0	1	0
Boxwood, hemlock, holly (English, American)			2	1	2	1
Douglas fir (seed tree)			25	15	20	15
Larch			5	3	10	4
Lilac (<i>Syringa</i> spp.)			3	1	3	1

	Sitka spruce (seed tree)	Late growth stage	25	15	25	15
	Spruce (seed tree and woodland)	Late growth stage	20	10	25	15
	Euonymus	Late growth stage	4	2	4	2
	Roses	Late growth stage	2	1	2	1
	Poinsettias	Late growth stage	2	1	2	1
	Blueberry (low and highbush)	Early growth stage	4	1	3	1
		Late growth stage	2	1	2	1
	Tree nuts (filberts, hazelnuts)	Early growth stage	15	5	20	10
		Late growth stage	5	3	10	4
	Peaches	Early growth stage	20	10	15	5
		Late growth stage	10	4	10	4
	Pears	Early growth stage	15	5	15	5
		Late growth stage	10	3	5	3
	Cherries (sour, sweet)	Early growth stage	10	4	10	3
		Late growth stage	5	2	4	2
Aerial	Flax, canary grass (seed production), peas (field and/or canning)	Fixed wing	1	0	1	0
		Rotary wing	1	0	1	0
	Cereals (barley, oats, rye, wheat), forage crops (clover, grasses, hay, grain), non-cropland (fence rows, field edges except adjacent to unregistered crops), pastures and wastelands, safflower, soybeans	Fixed wing	10	1	5	1
		Rotary wing	5	1	1	1
	Alfalfa (forage and seed production), clover (sweet, red, alsike)	Fixed wing	10	1	5	1
		Rotary wing	5	1	3	1
	Beans (pole, snap), soybeans	Fixed wing	5	1	5	0
		Rotary wing	3	1	1	0
	Hemlock	Fixed wing	90	0	150	15
		Rotary wing	55	0	90	1
	Blueberry (low and highbush)	Fixed wing	4	1	1	1
		Rotary wing	1	1	1	1

*For field sprayer application, buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.
When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that LAGON 480 E Insecticide contains a Group 1B insecticide/acaricide. Any insect/mite population may contain individuals naturally resistant to LAGON 480 E Insecticide and other Group 1B insecticide/acaricide. The resistant individuals may dominate the insect/mite population if this group of insecticides/acaricides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

2020-0144
2021-02-11

To delay insecticide/acaricide resistance:

- Where possible, rotate the use of LAGON 480 E Insecticide or other Group 1B insecticides/acaricides with different groups that control the same pests.
- Avoid application of more than the indicated number of sprays of LAGON 480 E Insecticide or other insecticides/acaricides in the same group in a season.
- Use tank mixtures with insecticides/acaricides from a different group when such use is permitted.
- Insecticide/acaricide use should be based on an IPM program that includes scouting, record keeping and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information and to report suspected resistance, contact the Technical Service, Loveland Products Canada Inc., 1-800-328-4678 or at www.uap.ca.

FOR GROUND APPLICATION: dilute with water to 200 - 300 L/ha unless otherwise stated.

FOR AERIAL APPLICATION: unless otherwise stated, dilute with water to 11 - 22 L/ha for blueberries, cereals (barley, oats, wheat, rye) and peas. Dilute with water to 10-30 L/ha for other field and vegetable crops when aerial application is specified.

AERIAL APPLICATION INSTRUCTIONS

Use only where aerial application is indicated on this label.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product.

Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the *Federal/Provincial/Territorial Committee on Pest Management and Pesticides*.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist).

Use low nozzle pressure (Below 300 kPa). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray in winds exceeding 10 - 15 km per hour. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat. Do not allow spray drift onto sensitive areas such as water, urban and residential areas.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) fine classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT direct spray above plants to be treated.

Turn off outward pointing nozzles at row ends, and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

Aerial application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) fine classification. Reduce drift caused by turbulent wingtip vortices. The nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Product Specific Precautions

If you have questions, call the manufacturer at 1-800-328-4678 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

2020-0144
2021-02-11

Volume: Apply the recommended rate in a minimum spray volume of 10 litres per hectare unless otherwise stated.

FIELD AND VEGETABLE CROPS:

Sprays should be applied when the pests first appear and repeated as the infestation warrants, or as otherwise directed. Applications should be made in sufficient water to obtain good coverage. Apply at the recommended rate; do not exceed a maximum spray volume of 1000 L per hectare unless otherwise stated. When aerial application is specified apply with aerial equipment and use 10 to 30L of water per hectare. Applications may be made by ground equipment unless otherwise specified.

FIELD AND VEGETABLE CROPS	PESTS CONTROLLED	AMOUNT PER HECTARE	AMOUNT PER 10 LITRES	PREHARVEST INTERVALS (DAYS)	MAXIMUM # OF APPLICATIONS PER YEAR
Alfalfa (for seed and forage production)	Aphids, Leafhoppers reduction of alfalfa weevil larvae	425 mL	4 mL	10	2
	COMMENTS: Do not graze or harvest for forage within 10 days after application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.				
	Blotch leafminers	550 mL	5 mL	10	1
COMMENTS: : Do not graze or harvest for forage within 10 days after application. Use a water volume of at least 200 L/ha with ground equipment for Blotch leafminers. Do not apply during bloom. May be applied by Ground and Aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
	Grasshoppers - nymphs	550 mL	5 mL	10	1
	Grasshoppers - adults or winged	850 - 900 mL	9 mL	10	2
	COMMENTS: Do not graze or harvest for forage within 10 days after treatment or when using a rate of 850 mL or higher: Do not graze or harvest for forage within 28 days after treatment. Follow provincial forecast. Apply when insects are present as young hoppers or signs of insect damage are evident. Apply when the grasshoppers are in the 2-4 nymphal stage. Best control will be achieved when application is made prior to wing development. Under severe insect pressure application should also be made to a 15 metre strip along fence rows around the field. DO NOT APPLY TO ADJACENT UNREGISTERED CROPS. The higher rate should be used when the proportion of mature and late nymphal stages in the population are high and spray penetration is inhibited by dense crop canopy. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.				
Alfalfa (seed production)	Lygus bugs Plant bugs	1.1 L	10 mL	28	1
COMMENTS: Apply just before alfalfa begins to bloom. Allow at least 10 days before placing leaf cutting bees in the field. Do not graze or harvest for forage with 28 days after treatment. May be applied by Ground and Aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
Asparagus	Asparagus aphid	2.3 L	35 mL	Not Applicable	2
COMMENTS: Ground application only. Do not apply during the crop bloom period or during the 5-day period before the crop blooms. For mature asparagus, applications can be made post-harvest only, until plant defoliation.					

2020-0144
2021-02-11

For immature asparagus, begin application mid-May. Apply by boom or field sprayer using 675 L of water per hectare. For immature asparagus; **DO NOT** harvest for feed or food. Ground application ONLY. Max number of applications per year is 2.

Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

Barley, Oats, Wheat (ALSO SEE CEREALS below)	Thrips	1 L	10 mL	35	2
COMMENTS: Ground Application: Use sufficient water to obtain good coverage (usually 110 – 275 L/ha). Same limitations apply for harvesting and grazing as in wheat section of this label. Ground and Aerial application. With aerial equipment use 10 to 30 L of water per hectare. Maximum number of applications per year is 2. Minimum application interval is 7 days. All activities: REI is 12 hours. Pre-grazing Interval (PGI): 14 days.					
Beans (Pole, Snap)	Aphids Grasshoppers Leafhoppers Leafminers Mexican bean beetles Mites, Lygus bugs Tarnished plant bugs	700 mL - 1 L	7 - 10 mL	7	2
COMMENTS: Use sufficient water for good coverage. Repeat applications as necessary. Do not feed or graze treated foliage to livestock. May be applied by Ground and Aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.					
Beets	Tarnished plant bugs	700 mL	7 mL	12	2
COMMENTS: Ground application ONLY. Minimum application interval is 7 days. All activities: REI is 3 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.					
Cauliflower Broccoli Brussels Sprouts	Aphids	700 mL - 1 L	7 - 10 mL	7 Broccoli Cauliflower 21 Brussels Sprouts	2
COMMENTS: To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period. Use sufficient water for good coverage. Repeat applications as necessary. Ground Application ONLY. Minimum application interval is 7 days. For all activities: REI is 5 days.					
Cauliflower Broccoli	Thrips	550 mL - 1.25 L	6 - 12 mL	7	2
To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period. Apply at 7 day intervals starting at head formation. Ground Application ONLY. Minimum application interval is 7 days. All activities: REI is 5 days.					
Canary Grass (Grown for seed)	Aphids	500 mL	5 mL	21	2
COMMENTS: Apply in aerial or ground application when there are more than 50 aphids per canary seed head between heading and soft dough stage. May be applied by Ground and Aerial application. May be applied a maximum of 2 times per year. Minimum application interval is 30 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when					

2020-0144
2021-02-11

most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Canola (rape)	Aphids Leafhoppers Grasshoppers	850 - 900 mL	9 mL	21	2
	COMMENTS: May be applied by ground and aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.				
Celery	Aphids-including carrot green peach foxglove aphids	700 mL	7 mL	7	2
	COMMENTS: Begin application 3 weeks after transplanting. Ground application ONLY. Minimum application interval is 7 days. All Activities, REI is 3 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.				
Cereals (barley, oats, wheat, rye)	Grasshoppers -nymphs, Say stink bugs	550 mL	5 mL	35	2
	Grasshoppers -adults	850 mL - 1 L	See comments		2
	COMMENTS: Do not graze within 14 days after treatment. Do not harvest for forage within 35 days after treatment. Follow provincial forecast. Apply when the grasshoppers are in the 2-4 nymphal stage. Best control will be achieved when application is made prior to wing development. Under severe insect pressure application should also be made to a 15 metre strip along fence rows around the field. DO NOT APPLY TO ADJACENT UNREGISTERED CROPS. The higher rate should be used when the proportion of mature and late nymphal stages in the population are high and spray penetration is inhibited by dense crop canopy. May be applied by Ground and Aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours.				
Chicory, Chinese Broccoli Bok Choi	Aphids	600 mL - 1 L	6 - 10 mL	7	2
	COMMENTS: Apply in sufficient water for thorough coverage. Use ground application to foliage. Apply when aphids are present. Ground application ONLY. Minimum application interval is 7 days. All activities: REI is 4 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.				
Eggplant	Tarnished plant bug	500 - 700 mL	5-7 mL	7	2
	COMMENTS: Apply when bugs are found and flowering is seen. Treat again in 7-10 days if bugs are found. Consult local authorities for proper timing. Do not apply when bees are foraging. Ground application ONLY. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.				
Flax	Potato Aphid	427 mL	4 mL	21	1
	COMMENTS: Apply from late flowering to early green bole stage; in sufficient water to provide good coverage. May be applied by either ground or aerial methods. Use in 200 L/ha of water for ground and 11 - 22 L of water/ha for aerial equipment. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.				
Forage crops	Aphids	425 - 550 mL	4 - 5 mL	2	2

2020-0144
2021-02-11

(grasses, alfalfa,
hay, clover)

(Only suppression of Russian Wheat aphid is provided)

Grasshoppers- low infestations	425 – 550 mL		2	2
nymphs	550 mL		2	2
adults	850 mL – 1L		28	2
Lygus Bug	425 mL		2	2
Plant Bug	425 mL		2	2

COMMENTS: TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms. Do not graze or harvest for forage within 2 days after treatment at the 425 or 550 mL rate. Do not graze or harvest for forage within 28 days after treatment at the 850 mL – 1 L rate. May be applied by either ground or aerial methods. Minimum application interval is 7 days. All activities: REI is 12 hours.

Leafy Vegetables: Lettuce, Swiss chard, Beet greens and Turnip greens	Aphids Leafhoppers	700 mL	7 mL	14	2
COMMENTS: Ground application ONLY. Use sufficient water for good coverage. Do not use on Head Lettuce. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period. Minimum application interval is 7 days. Interval between last application and harvest:					
	Lettuce	14 days	Beet greens	21 days	
	Swiss chard	14 days	Turnip greens	14 days	
Beet Greens, Turnip Greens and Lettuce: All Activities, REI is 3 days.					

Kale	Aphids Leafhoppers	700 mL	7 mL	14	2
COMMENTS: Ground application ONLY. Use sufficient water for good coverage. May be applied a maximum of 2 times per year. Minimum application interval is 15 days. Interval between last application and harvest: 14 days. All Activities, REI is 3 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.					

Peas (canning & field)	Aphids	275 - 380 mL	2.5 - 3.6 mL	3	2
COMMENTS: Apply by aircraft or by ground equipment. Use sufficient water for good coverage. Repeat application as necessary. Do not feed or graze for forage within 21 days after last application. Dilute for ground (100-300 L of water/ha) and aerial application (10-30 L of water/ha). Minimum application interval is 14 days. Maximum application rate is 0.18 kg a.i./ha. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.					

Peppers	Aphids, Maggots	700 mL - 1L	7 - 10 mL	3	2
COMMENTS: Use sufficient water for good coverage. Repeat application as necessary. Ground application ONLY. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.					

Potatoes, Tomatoes (field)	Aphids Leafhoppers Tarnished plant bugs	550 mL - 1.0 L	5 - 9 mL	7	2
COMMENTS: Use sufficient water for good coverage. Do not apply more than 2 applications per season. Consult local agricultural authorities for proper timing. Ground application ONLY. Minimum application interval is 7 days. Maximum application rate is 0.48 kg a.i./ha All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions					

2020-0144
2021-02-11

section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Pastures Wastelands	Grasshoppers -nymphs -adults	550 mL 850 mL-1 L		2 See Comments	2
<p>COMMENTS: Follow provincial forecast. Apply when insects are present as young hoppers or signs of insect damage are evident. Apply when the grasshoppers are in the 2-4 nymphal stage. Best control will be achieved when application is made prior to wing development. Under severe insect pressure application should also be made to a 15 metre strip along fence rows around the field. DO NOT APPLY TO ADJACENT UNREGISTERED CROPS. The higher rate should be used when the proportion of mature and late nymphal stages in the population are high and spray penetration is inhibited by dense crop canopy. Use high rate for adult grasshopper control on wasteland. Remove cattle before spraying. Do not graze or harvest for forage within 2 days after treatment at the 550 mL rate. Do not graze or harvest for forage within 7 days after treatment at the 850 mL rate. Do not graze or harvest for forage within 28 days at the 1L rate. May be applied by ground and aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>					
Safflower	Grasshoppers (nymphs and adults)	550 mL-1L		21	2
<p>COMMENTS: Apply when damage is apparent and more than 15 grasshoppers per square metre are found in the crop. Repeat the application only when necessary. Use sufficient water for good coverage. May be applied by ground and aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.</p>					
Soybeans	Aphids bean beetles leafhoppers tarnished plant bugs, lygus bugs	0.7-1 L	7-10 mL	30	2
<p>COMMENTS: Use sufficient water for good coverage. Repeat applications as necessary. Do not exceed 3 applications per season. Do not feed or graze treated foliage to livestock. May be applied by Ground and Aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>					
	Spider mites (Two-spotted)	1 L	10 mL	30	2
<p>COMMENTS: Apply by ground application in sufficient water for thorough coverage. Do not feed or allow livestock to graze treated forage. May be applied by ground and aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>					
Sweet Clover Red Clover Alsike Clover	Sweet clover weevil	850 mL - 1.1 L		28	2
<p>COMMENTS: Do not graze or harvest for forage within 28 days after treatment. May be applied by ground and aerial application. Minimum application interval is 7 days. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.</p>					
Wheat	Orange wheat,	1 L		21	2

2020-0144
2021-02-11

(ALSO SEE
CEREALS above)

blossom midge,
Russian wheat
aphid

COMMENTS: If midge population persists at 7 days apply a second treatment. Use a water volume of at least 100 L/ha with ground equipment and 50 L/ha by air. Higher volume will provide more thorough coverage. If adult midges are present (1 midge/4-5 wheat heads), sprays should be applied when 25% of the wheat head has fully emerged from the boot but before flowering has begun. At this stage, wheat first becomes susceptible to attack by egg-laying females. Applications should be made in the late afternoon or evening when temperatures exceed 15°C and the wind speed is less than 10 km/h. Higher volume sprays will improve penetration of the crop. Proper timing of application is essential for control. Do not harvest within 21 days of the application. Do not apply to areas where bees are actively foraging or near apiaries as product is toxic to bees. Do not graze or use for forage within 7 days of application. Consult provincial authorities for further information concerning rates and time of application. Minimum application interval is 7 days. All activities: REI is 12 hours.

FRUIT CROPS:

APPLY WITH GROUND EQUIPMENT ONLY UNLESS OTHERWISE NOTED. FOR DILUTE GROUND APPLICATIONS, Apply at the recommended rate; do not exceed a maximum spray volume of 1000 L per hectare unless otherwise stated. For Cherry, Peach and Pear, the rate per hectare is based on using 3000 L of spray on full sized trees, 4.5-5.5 metres high, Apply at the recommended rate; do not exceed a maximum spray volume of 3000 L per hectare unless otherwise stated. Trees smaller or larger may require more or less LAGON 480 E. Follow provincial calendar guidelines as to water volumes for the size of tree being sprayed.

FRUIT TREE CROPS	PEST CONTROLLED	AMOUNT PER HECTARE	AMOUNT PER 10 L	AMOUNT PER 1000 L	PREHARVEST INTERVALS	MAXIMUM # OF APPLICATIONS PER YEAR
Pears	Tarnished plant bugs		6.25 mL	625 mL	28	2
	COMMENTS: Pears: Apply at pre-bloom. The minimum application interval is 10 days. Maximum rate 1.44 kg a.i./ha. Thinning: REI is 28 days. Hand harvesting, hand-line irrigation: REI is 17 days. All other activities: REI is 1 day. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
			10 – 12.5 mL	1 – 1.25 L	28	2
	Pear Psylla					
	COMMENTS: Use sufficient water for good cover spray. The minimum application interval is 10 days. Maximum application rate is 1.44 kg a.i./ha. Thinning: REI is 28 days. Hand harvesting, hand-line irrigation: REI is 17 days. All other activities: REI is 1 day. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
	Aphids		10 – 12.5 mL	1 – 1.25 L	28	2
	COMMENTS: Use sufficient water for good cover spray. The minimum application interval is 10 days. Maximum application rate is 1.44 kg a.i./ha. Thinning: REI is 28 days. Hand harvesting, hand-line irrigation: REI is 17 days. All other activities: REI is 1 day. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
	Mites		10 – 12.5 mL	1 – 1.25 L	28	2
	COMMENTS: Use sufficient water for good cover spray. The minimum application interval is 10 days. Maximum application rate is 1.44 kg a.i./ha. Thinning: REI is 28 days. Hand harvesting, hand-line irrigation: REI is 17 days. All other activities: REI is 1 day. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					
Peaches (non-bearing)	Aphids, Mites		10 -15 mL	1 – 1.5 L	40	2
	Tarnished plant bugs	1.7 L	5 mL	500 mL	40	2
	COMMENTS: Spray when insects first appear and repeat as necessary using sufficient water for good coverage. Some defoliation may occur under cold wet soil and/or slow drying conditions during and following application. Least injury has occurred when sprays were applied in the morning while temperatures are moderate and drying conditions are fairly good. The minimum application interval is 10 days. Thinning: REI is 32 days. Hand harvesting, hand-line irrigation: REI is 20 days. All other activities: REI is 3 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.					

2020-0144
2021-02-11

Sour Cherries	Cherry fruit fly Maggots	2.25 L	5 mL	500 mL	21	2
<p>COMMENTS: Apply no later than 6 days after the first adult fly has been trapped and repeat in 14 days if required. Thinning: REI is 20 days. Hand harvesting, hand-line irrigation: REI is 9 days. All other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.</p>						
Sweet Cherries	Cherry fruit fly maggots	2.25 L	5 mL	500 mL	21	2
<p>COMMENTS: Apply no later than 6 days after the first adult fly has been trapped and repeat in 14 days if required. Thinning: REI is 20 days. Hand harvesting, hand-line irrigation: REI is 9 days. All other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.</p>						
SMALL FRUIT CROPS	PESTS CONTROLLED	AMOUNT PER HECTARE	AMOUNT PER 10 LITRES	AMOUNT PER 1000 LITRES	PREHARVEST INTERVALS (DAYS)	MAXIMUM # OF APPLICATIONS PER YEAR
Blueberry Lowbush Highbush	Blueberry maggot	580 – 830 mL 830 mL	6 – 8 mL		21	2
<p>COMMENTS: Apply pesticide in sufficient water for good coverage, one week after first berries ripen. Use no more than 1000 L spray/ha. Make 1 or 2 applications between mid to late July. May vary depending on area and season. May be applied with ground equipment or aerial equipment. Hand harvesting (high bush): REI is 9 days. Hand harvesting (low bush) and all other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.</p>						
Strawberries (bearing)	Tarnished plant bug	2.75 L	12.5 mL	1.25 L	7	2
<p>COMMENTS: Apply first spray when first blooms appear and the second application 10-12 days later if needed. May be applied a maximum of 2 times per year. Minimum application interval is 10 days. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>						
Strawberries (bearing & non-bearing)	Aphids, mites	2.25 L	10 mL	1 L	7	2
<p>COMMENTS: Use sufficient water for good cover spray. Spray when insects first appear and repeat at monthly intervals as required. May be applied a maximum of 2 times per year. Minimum application interval is 10 days. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.</p>						
CROPS	PEST	RATE PER HECTARE		REMARKS		
Filberts, Hazelnuts	Aphids	5L		<p>Use one application per season. Apply when aphids appear. DO NOT USE WITHIN 45 DAYS OF HARVEST. Primarily for use on young plantings Thinning: REI is 34 days. Hand harvesting, hand-line irrigation: REI is 21 days. All other activities: REI is 5 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict</p>		

applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

ORNAMENTALS

GENERAL COMMENTS ORNAMENTAL PLANTS

APPLY ONLY WITH GROUND APPLICATION EQUIPMENT

1. LAGON 480 E is a true systemic insecticide. A systemic insecticide is absorbed into the system of the plant upon application, and as with all systemic materials, may cause reactions in specific plants which are not predictable nor common to all members of the species. Do not over-dose or over-spray.
2. FOR PROPER TIMING of treatments for the control of specific pests on ornamental plants, consult local agricultural authorities. In general, apply adequate spray for good coverage when pests appear or when damage is first observed, unless otherwise indicated.
3. EXPERIENCE has shown when dealing with species that may be injured by foliar application, soil drenches are safer on those plants for which they are recommended. When using foliar sprays, do not apply during heat of the day or when temperatures are excessively high.
4. DO NOT USE on ornamental plants not listed.
5. WHEN IN DOUBT about plant safety, it is advisable to spray a single branch or dip several leaves in the spray solution. A waiting period of 3 or 4 days will usually indicate if plant damage will result.
6. DO NOT SPRAY the following species unless you have tested single branches or leaves to ascertain safety:

American Elm	Mop-head Elm
Norway Maple	Mock Orange
Sunburst Locust	Morraine Locust

 or other ornamentals not listed in this pamphlet.
7. There has been no report of damage by LAGON 480 E to any of the common evergreen species grown in Canada provided recommended rates have been followed.
8. Do not apply foliar spray during the heat of the day or when temperatures are exceedingly high.
9. Apply at the recommended rate; do not exceed a maximum spray volume of 1000 L per hectare unless otherwise stated.

OUTDOOR FLOWERING PLANTS	PESTS CONTROLLED	AMOUNTS PER 1000 L OF WATER	AMOUNTS PER 10 L OF WATER
Azaleas	Lace bugs, Leafminers, Mites Whiteflies COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL
Camellias	Aphids, Mites, Camellia scales FOLIAR SPRAYS: Apply 2 sprays 6 weeks apart the first year followed by annual applications soon after first growth begins in the spring. The minimum application interval is 10 days. May be applied a maximum of 2 times per year. All Activities, REI is 2 days. SOIL DRENCH: Apply LAGON 480 E as a soil drench around the base of plants in early spring at the rate of 60 mL per 10 L of water per plant up to 2 metres tall. Increase this rate proportionately for larger plants. Soil drench: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10mL
Day Lilies	Aphids, Thrips COMMENTS: May be applied a maximum of 1 time per year. All activities: REI 7 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	2 L	20 mL
Euonymus	Aphids, Scales COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	2 L	20 mL

2020-0144
2021-02-11

Gardenias	Tea scales, White flies COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL
Gerberas	Thrips COMMENTS: May be applied a maximum of 1 time per year. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL
Gladiolus	Aphids, Thrips COMMENTS: May be applied a maximum of 1 time per year. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL
Iris	Aphids, Iris Borers, Thrips For borer control spray when new leaves are 12 to 15 cm tall. COMMENTS: May be applied a maximum of 1 time per year. All activities: REI is 7 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	2 L	20 mL
Poinsettias (outdoor plants)	Aphids, White flies, Mites, Mealy bugs COMMENTS: May be applied a maximum of 1 time per year. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL
Roses (outdoor plants)	Aphids, Leafhoppers, Mites, Thrips Apply thoroughly during growing season in a spray. The minimum application interval is 28 days. May be applied a maximum of 2 times per year. All Activities, REI is 2 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms..	1 L	10 mL

**OUTDOOR
ORNAMENTALS
AND CHRISTMAS
TREES**

	PESTS CONTROLLED	AMOUNTS PER 1000 L OF WATER	AMOUNTS PER 10 L OF WATER
Christmas Trees: Balsam Fir, Spruce, Hemlock, Pine (Mugho, Red, Scots)	Aphids, mites, Scales, Spruce needle miners, Spruce budworms COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 18 days. Hand line irrigation: REI is 3 days. All other activities: 12 hours.	1.5 L	15 mL
	Bagworms, European pine shoot moths, Nantucket pine tip moths, Zimmerman pine moths COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 18 days. Hand line irrigation: REI is 3 days. All other activities: REI is 12 hours.	2 L	20 mL
Arborvitae	Aphids, Bagworms, Mites COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All activities: REI is 12 hours.	2 L	20 mL
Birch	Aphids, Birch Leafminers FOLIAR SPRAY: For leaf miners apply when leaves are expanded (about mid-May); repeat the last week in June. For all activities: REI is 12 hours. SOIL DRENCH: For full-season control LAGON 480 E may be used as a soil drench once in early May when the leaves start to open. Apply LAGON 480 E undiluted at a rate of 11 mL/cm of stem basal diameter (cumulative total of all stems) evenly into small shallow holes made with a sharp instrument at the dripline of the tree or clump. Close the holes with soil. Drench thoroughly with water	500 mL	5 mL

immediately so as to saturate the root system. One soil drench treatment in May should give control of birch leafminers for the complete season. Do not make more than one soil drench treatment per season. Soil drenches may not be effective on newly-transplanted birches where the root system is not well established. Some hybrid varieties eg: cutleaf and silver birch may be damaged by excessive amounts of LAGON. Therefore, apply no more than the recommended dosage. Soil drench: REI is 12 hours.

COMMENTS: To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.

Boxwood	Mites, Leafminers, Mealy bugs	1 L	10 mL
	COMMENTS: For leafminers, apply in spring when leafminer flies first appear or in early summer for control of larvae in the infested leaves. The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 7 days. All other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.		
Cedar	Mites	2 L	20 mL
	COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 18 days. Hand-line irrigation: REI is 3 days. All other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.		
Douglas-Fir (seed tree)	Cone insects (cone or gall midges, cone moths, seed chalcids, scale midges). 2 L per 100 litres of water. Maximum rate 20 L of product (= 9.6 kg a.i.) per hectare. Application should be made when cones are at or near the pendant stage. Complete coverage of the cones and foliage in the cone-bearing area of the tree is important for satisfactory results. Maximum of two applications per year. Minimum application interval: 10 days Seed cone harvesting: 48 days REI. Scouting: 5 days REI. Grading, animal control, baiting: 1 day REI.		
Hemlock	Mites, Scales, Spruce needle miners Spruce budworms	1 L	10 mL
	COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 13 days. All other activities: REI is 12 hours.		
Holly (English & American)	Mites, Soft Scale, Leafminers	1 L	10 mL
	COMMENTS: For leafminers apply in spring when leafminer flies first appear, or in early summer for the control of larvae in the infested leaves. The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All activities: REI is 12 days. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.		
Juniper	Aphids, Bagworms, Midges, Mites	2 L	20 mL
	COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. All activities: REI is 12 hours.		
Larch	Cone Maggots, Seed Wasps	5 L	50 mL
	COMMENTS: One application per season; apply within 3 weeks after pollination; spray to runoff on foliage; ground hydraulic sprayer or mist sprayer. All Activities, REI is 1 day. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.		
Lilacs (Syringa spp.)	Leafminers	1.25 L	12.5 mL
	COMMENTS: Apply two sprays 6 weeks apart. Application should be made as soon as any signs of the leafminer appear, or, first application should be started in early June. The minimum application interval is 10 days. May be applied a maximum of 2 times per year. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms. All activities: REI is 15 days.		
Oak	Golden oak scales	2 L	20 mL
	COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 18 days. Hand-line irrigation: REI is 3 days. All other activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. DO NOT apply during the crop blooming period or during the 5-day period before the crop blooms.		

2020-0144
2021-02-11

Pine (Mugho, Red, Scots)	Aphids, Bagworms, European pine shoot moths, Nantucket pine tip moths, Zimmerman pine moths	2 L	20 mL
	Red and black-headed pine sawfly larvae	1 L	10 mL

COMMENTS: For pine shoot moths apply early in the spring and again in mid-June, thoroughly wetting branch tips. COMMENTS: The minimum application interval is 14 days. May be applied a maximum of 2 times per year. Thinning: REI is 18 days. Hand line irrigation: REI is 3 days. All other activities: REI is 12 hours.

Sitka spruce (seed tree): Sitka spruce weevils. 20.8 L per 1000 litres of water. Maximum rate 20.8 L of product (= 10 kg a.i.) per hectare. Thoroughly spray the terminal growth to the point of runoff at the time of egg laying (usually during the first half of May). Maximum two applications per year. Minimum application interval: 10 days. Seed cone harvesting: 49 days REI. Scouting: 5 days REI. Hand-pruning, staking, tying: 27 days REI. Grading, animal control, baiting: 1 day REI.

Spruce (seed tree and woodland use): Seed and cone insects. 10.4 - 20.8 L per 1000 L water. Maximum rate 20.8 L of product (10 kg a.i) per hectare. Thoroughly spray cone-bearing portions of tree to point of runoff when strobili cones are closed, and turning, but before they reach the horizontal position. Contact your local forest pest control office for more information on timing, and spray application, as well as spruce species that may be treated. Maximum of two application per year. Minimum application interval: 10 days. Seed cone harvesting: 49 days REI. Scouting: 5 days REI. Hand pruning, staking, tying: 27 days REI. Grading, animal control, baiting: 2 days REI.

Taxus	Mites, Fletcher scale, Mealy bugs	2 L	20 mL
-------	-----------------------------------	-----	-------

COMMENTS: For Fletcher scale, spray at the end of June and again 14 days later when crawlers are moving. May be applied a maximum of 2 times per year. Minimum application interval is 14 days. All activities: REI is 12 hours.

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than Loveland Products Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Loveland Products Canada Inc. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crop listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Loveland Products Canada Inc. harmless from any claims based on efficacy and/or phytotoxicity in connection with the use described below.

Highbush Blueberry

Crop	Pest	Rate per 1000 L water	Maximum No. of Applications	Reapplication Interval
Highbush Blueberry	Spotted Wing Drosophila	830 mL	2	15 days

COMMENTS: Apply post-harvest only, to control spotted wing Drosophila (adults and larvae in fruit) that may otherwise infest adjacent crops. Timing varies, depending on variety of highbush blueberry. Foliar application only, using conventional ground application equipment. Use sufficient water volume to ensure thorough coverage, to a maximum of 1000 L/ha. REI is 12 days.

2020-0144
2021-02-11

RESTRICTED USES - FORESTRY USE

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Product Act to use this product in a way that is inconsistent with the directions on the label.
CAUTION: Forest Pest Control - birds and other wildlife in treated area may be harmed.
Apply at the recommended rate; do not exceed a maximum spray volume of 1000 L per hectare unless otherwise stated.

RESTRICTED USE - AERIAL APPLICATION

TREE	PESTS CONTROLLED	COMMENTS
Hemlock, Spruce, Balsam fir	Mites, Scales, Spruce needle miners, Spruce budworms	For aircraft application use 1.5 L of LAGON 480 E in 5-20 L of water/ha, applied as a uniform spray of fine droplets. Application should be made as soon as the larvae hatch. COMMENTS: May be applied a maximum of 1 time per year. Thinning: REI is 13 days. All other activities: REI is 12 hours.

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the use described below were developed by persons other than Loveland Products Canada Inc. and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Loveland Products Canada Inc. itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crop listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Loveland Products Canada Inc. harmless from any claims based on efficacy and/or phytotoxicity in connection with the use described below.

ANNUAL CANARYGRASS

CROP	PEST	RATE PER HA	MAXIMUM NO. OF APPLICATIONS PER YEAR	PREHARVEST INTERVALS (DAYS)	AMOUNT PER 10 LITRES
Annual canarygrass	Aphids	500 mL	1	21	5 mL

COMMENTS: Apply in aerial or ground application when there are more than 50 aphids per canary seed head between heading and soft dough stage. May be applied by Ground and Aerial application.

May be applied a maximum of 1 time per year. All activities: REI is 12 hours. To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section. TOXIC to bees. Avoid application during the crop blooming period. If applications must be made during the crop blooming period, restrict applications to evening when most bees are not foraging. When using managed bees for pollination services, DO NOT apply during the crop blooming period.

Annual canarygrass seeds can be harvested for human consumption.

Annual canarygrass grown for human consumption is not to be cut for feed or grazed.

2020-0144
2021-02-11

DECONTAMINATION

If accidental spillage of LAGON 480 E Insecticide should occur, scrub contaminated area immediately with a strong laundry soap solution or use household lye. Detergents are not satisfactory for this purpose. Repeated scrubblings are necessary on plain wood surfaces. Avoid spillage on all types of floor tile or linoleum.

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Dimethoate is an organophosphate that is a cholinesterase inhibitor. Typical symptoms of overexposure to cholinesterase inhibitors include headache, nausea, dizziness, sweating, salivation, and runny nose and eyes. This may progress to muscle twitching, weakness, tremor, incoordination, vomiting, abdominal cramps and diarrhea in more serious poisonings. A life-threatening poisoning is signified by loss of consciousness, incontinence, convulsions and respiratory depression with a secondary cardiovascular component. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate degree of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as pralidoxime chloride, may be therapeutic if used early; however, use only in conjunction with atropine. In cases of severe acute poisoning, use antidotes immediately after establishing an open airway and respiration. With oral exposure, the decision of whether to induce vomiting or not should be made by an attending physician. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

STORAGE

Do not use, pour, spill, or store near heat, sparks, or open flame. Keep out of direct sunlight. Ship and store between 4°C and 30°C.

DISPOSAL

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.