



## **PRECAUTIONS**

**HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN**

**EYE IRRITANT**

**KEEP OUT OF REACH OF CHILDREN**

1. DO NOT APPLY BY AIR.
2. Do not take internally. Handle with care and mix only in a closed container. Use only in a well ventilated area.
3. May cause irritation of eyes and skin. Avoid inhalation of vapour or spray mist and contact with eyes, skin or clothing.
4. Wash thoroughly after handling and before eating, drinking, smoking or using the bathroom. Take a shower immediately after work.
5. If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.
6. Do not contaminate domestic or irrigation water, lakes, ditches, streams or ponds by the cleaning of equipment or the disposal of wastes.
7. Do not apply where run-off or erosion is likely to occur.
8. **DO NOT** use in residential areas. Residential areas are defined as any use site where bystanders or the public, including children, could be exposed during or after application. This includes areas in and around homes, schools, public buildings or any other areas where they could be exposed.

## **PROTECTIVE CLOTHING AND EQUIPMENT**

### **All Application Equipment**

Wear a hat, a long-sleeved shirt, long pants, chemical-resistant gloves, socks and rubber boots during mixing, loading, application, clean-up and repair. In addition, wear coveralls and protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

## **PHYSICAL OR CHEMICAL HAZARDS**

**COMBUSTIBLE.** Do not use or store near heat or open flame.

## **FIRST AID**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

## **TOXICOLOGICAL INFORMATION**

The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or esophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient. High concentrations of MCPA may cause severe irritation to the eyes. Symptoms of overexposure to MCPA could include slurred speech, twitching, jerking and spasms, drooling, low blood pressure and unconsciousness. Treat symptomatically.

## **ENVIRONMENTAL PRECAUTIONS**

This product contains aromatic petroleum distillates, which are toxic to aquatic organisms. Avoid contamination of aquatic systems during application. DO NOT contaminate these systems through direct application, disposal of waste or cleaning equipment.

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to, heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted, fine-textured, or low in organic matter such as clay).

Avoid applying this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

## **STORAGE**

Store in original, tightly-closed container and DO NOT allow water to be introduced into this container. DO NOT ship or store near food, feed, seed and fertilizers. Store in cool (above 5°C) dry, locked, well ventilated area without floor drain. If product is frozen, bring to room temperature and agitate before use. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.

## **DISPOSAL**

### **Recyclable Containers:**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

### **Returnable Containers:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**Refillable Containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label.



## **PRECAUTIONS**

**HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN**

**EYE IRRITANT**

**KEEP OUT OF REACH OF CHILDREN**

1. DO NOT apply by air.
2. Do not take internally. Handle with care and mix only in a closed container. Use only in a well ventilated area.
3. May cause irritation of eyes and skin. Avoid inhalation of vapour or spray mist and contact with eyes, skin or clothing.
4. Wash thoroughly after handling and before eating, drinking, smoking or using the bathroom. Take a shower immediately after work.
5. If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.
6. Do not contaminate domestic or irrigation water, lakes, ditches, streams or ponds by the cleaning of equipment or the disposal of wastes.
7. Do not apply where run-off or erosion is likely to occur.
8. **DO NOT** use in residential areas. Residential areas are defined as any use site where bystanders or the public, including children, could be exposed during or after application. This includes areas in and around homes, schools, public buildings or any other areas where they could be exposed.

## **PROTECTIVE CLOTHING AND EQUIPMENT**

### **All Application Equipment**

- Wear a hat, a long-sleeved shirt, long pants, chemical-resistant gloves, socks and rubber boots during mixing, loading, application, clean-up and repair. In addition, wear coveralls and protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

## **PHYSICAL OR CHEMICAL HAZARDS**

**COMBUSTIBLE.** Do not use or store near heat or open flame.

## **FIRST AID**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

## **TOXICOLOGICAL INFORMATION**

The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or esophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient. High concentrations of MCPA may cause severe irritation to the eyes. Symptoms of overexposure to MCPA could include slurred speech, twitching, jerking and spasms, drooling, low blood pressure and unconsciousness. Treat symptomatically.

## **ENVIRONMENTAL PRECAUTIONS**

This product contains aromatic petroleum distillates, which are toxic to aquatic organisms. Avoid contamination of aquatic systems during application. DO NOT contaminate these systems through direct application, disposal of waste or cleaning equipment.

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to, heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted, fine-textured, or low in organic matter such as clay).

Avoid applying this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sand, loamy sand and sandy loam soils) and/or the depth to the water table is shallow.

## **STORAGE**

Store in original, tightly-closed container and do not allow water to be introduced into this container. DO NOT ship or store near food, feed, seed and fertilizers.

Store in cool (above 5°C) dry, locked, well-ventilated area without floor drain. If product is frozen, bring to room temperature and agitate before use. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.

## **DISPOSAL**

### **Recyclable Containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

### **Returnable Containers:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

### **Refillable Containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

### GENERAL INFORMATION

Certain Herbicide should be diluted in water and applied by ground application equipment for the control of the following broadleaved weeds:

1.5 L/ha (low rate)

Canada thistle♦ (low infestations)	prickly lettuce
burdock	radish, wild
cocklebur	ragweed
flixweed (spring rosettes 2-4 leaf)	shepherd's-purse (spring rosettes only)
♦♦field horsetail	stinkweed (spring rosettes)
lamb's-quarters	sunflower, annual
mustard, wild	sunflower, volunteer
♦♦plantain	vetch

♦Season long control, with some regrowth in the fall.

♦♦Top growth only

2.0 L/ha (high rate) - the above listed weeds plus

buckwheat, tartary	pigweed, redroot
buckwheat, wild	pigweed, Russian
Canada thistle♦ (medium to high infestations)	scentless chamomile (2 to 4 leaf)
canola, volunteer	smartweed
common groundsel	sow-thistle, annual
dandelion (spring rosettes only)	sow-thistle, perennial (top growth)
kochia (suppression) (2 to 4 leaf)	

### DIRECTIONS FOR USE

#### FIELD SPRAYER APPLICATION

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Best results are obtained when Certain Herbicide is applied to actively growing weeds in the 1 to 4 leaf (seedling) stage and when Canada thistle is between 10 cm and up to but not including the early bud stage. Poor weed control may result under cool or dry conditions.

#### Wheat, Barley, Oats

Apply to the crop from 3 leaf to just before flag leaf. Apply Certain Herbicide as a ground treatment using enough spray volume to provide adequate spray coverage. Apply 100 to 150 L/ha total spray volume by ground. Higher spray volumes will provide better coverage and weed control, particularly when the weeds are not actively growing. Maximum 1 application per year.



**Flax (including low linolenic acid varieties)**

Apply to the flax when the crop is 5 to 15 cm tall. Early spraying will reduce the risk of crop injury. Apply Certain Herbicide in 100 to 150 L/ha total spray volume by ground equipment at spray pressures of 200 to 275 kPa. For optimum spray coverage, use flat fan nozzles tilted 45° forward to ensure better coverage. Maximum 1 application per year.

**NOTE:** Some delay in maturity and crop injury may result in flax from the use of the high rate but any potential yield reductions will usually be more than offset by increased yields due to better weed control.

**Spray Preparation**

Add half the required amount of water to the spray tank, then add with agitation the required amount of Certain Herbicide and the tank-mix partner if applicable. Add the balance of the water and continue to agitate while spraying.

The use of flood jet or hollow cone nozzles is not recommended because of uneven and inadequate spray coverage. Ensure thorough and uniform spray coverage over the entire leaf area of the target weeds.

**GENERAL USE PRECAUTIONS****DO NOT APPLY BY AIR.**

DO NOT use in greenhouses.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

**Sensitive Plants**

Do not apply Certain Herbicide directly to or otherwise permit it to come into contact with sunflowers, legumes, fruit or vegetable crops, flowers or other desirable broadleaved plants and take precautions to avoid spray mists drifting onto them.

**Crop Rotation**

Fields previously treated with Certain Herbicide can be seeded to wheat, barley, oats and rye (not underseeded to forage legumes, clover or alfalfa), canola, corn, field peas, flax, forage grasses, mustard, sugar beets or can be summer-fallowed. Seed only those crops listed above in the year following treatment. Do not seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application, delay seeding field peas an additional 12 months (total 22 months following application).

**Preharvest/Grazing Interval**

- Do not cut or graze treated fields of wheat, barley, oats or flax within 7 days after application.
- Do not harvest wheat, barley, oats or flax within 60 days after application, or within 7 days after application when harvesting for forage.
- Withdraw meat animals from treated fields at least 3 days before slaughter.

**Stress Conditions**

When weeds are stressed due to drought, flooding or prolonged hot or cool temperatures (8°C or less), control can be reduced or delayed since weeds are not actively growing. Weed escapes may occur under

prolonged stress conditions or low fertility. Do not make applications to weeds stressed longer than 20 days due to lack of moisture as unsatisfactory control can result.

### To Reduce Spray Drift

1. Use coarse sprays at low pressures (200 to 275 kPa).
2. Use 100 to 200 L/ha of spray solution.
3. Spray when the wind velocity is 16 km/hr or less.

### Equipment Clean-Up

Equipment used to apply Certain Herbicide should not be used to apply other pesticides to sensitive crops without thorough cleaning.

### TANK MIXTURES

DO NOT mix or apply this product with any other additive, pesticide or fertilizer except as specifically recommended on this label.

### TANK-MIXES WITH CERTAIN HERBICIDE FOR BROADLEAVED WEED AND ANNUAL GRASS CONTROL IN WHEAT AND BARLEY

Certain Herbicide may be tank-mixed with one of the tank-mix partners indicated in the table below for wide spectrum control. All broadleaved weeds listed elsewhere on this label plus the annual grasses listed below for each tank-mix partner will be controlled by the tank-mix. Follow all precautions, directions, and recropping/rotation limitations that appear on both product labels and follow those that are the most restrictive.

Herbicide Tank-Mix Partner	Crops	Certain Herbicide + tank-mix partner rate product/ha	Annual Grasses	Adjuvant Rate
Achieve Liquid Herbicide	spring wheat, durum wheat, barley	1.5 - 2.0 L + 0.5 L	wild oats, green & yellow foxtail, barnyard grass, Persian darnel	Turbocharge: 0.5 % v/v
Assert 300 SC Herbicide	spring wheat, durum wheat, barley	1.5 - 2.0 L + 1.3 L or 1.5 - 2.0 L + 1.6 L	wild oats, 1 to 3 leaf or wild oats, 4 leaf	Refer to Assert 300 SC label
Axial 100 EC Herbicide	spring wheat, barley	1.5 - 2.0 L + 600 mL	wild oats, green foxtail, yellow foxtail, volunteer oats, volunteer canaryseed, proso millet	Adigor 700 mL/ha
Everest Solupak 70 DF	spring wheat	1.5 - 2.0 L + 43 g	wild oats, green foxtail	Refer to Everest Solupak 70 DF label

### TANK-MIXES WITH CERTAIN HERBICIDE FOR WIDE SPECTRUM BROADLEAVED WEED CONTROL

Certain Herbicide may be tank-mixed with Refine Extra Herbicide for wide spectrum broadleaved weed control. All broadleaved weeds listed elsewhere on this label plus those listed on the Refine Extra label will be controlled by this tank-mix. Follow all precautions, directions, and recropping/rotation limitations that appear on both product labels and follow those that are the most restrictive.

Herbicide Tank-Mix Partner	Crops	Certain Herbicide + Refine Herbicide product /ha	Extra rate	Additional Weeds	Adjuvant Rate
Refine Extra Herbicide	spring wheat, spring barley	1.5 L + 20 g		refer to the list of broadleaved weeds on the Refine Extra Herbicide label	Agral 90, AgSurf or Citowett Plus: 0.2 % v/v

**NOTE TO USER:** READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS: The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than SHARDA Cropchem Limited under the User Requested Minor Use Label Expansion program. For these uses, SHARDA Cropchem Limited has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

#### **DIRECTIONS FOR USE**

##### **ESTABLISHED TIMOTHY FOR HAY AND FORAGE PRODUCTION IN WESTERN CANADA**

For the control of labeled weeds on established timothy for hay and forage production in Western Canada, apply Certain Herbicide at a rate of 2 L/ha. Apply only one post-emergent application per season. DO NOT apply within 7 days of harvest. For use in the Prairie Provinces and the Peace River, Okanagan and Creston regions of British Columbia only.

##### **ESTABLISHED TIMOTHY FOR SEED PRODUCTION IN WESTERN CANADA**

For the control of labeled weeds on established timothy for seed production in Western Canada, apply Certain herbicide at a rate of 1.5 - 2 L/ha. Apply only one post-emergent application per season. Apply to the crop from 3 leaf to just before flag leaf. Do not cut or graze treated fields within 7 days after application. For use in the Prairie Provinces and the Peace River, Okanagan and Creston regions of British Columbia only. Timothy is particularly sensitive to stress conditions, and some crop injury or seed yield reduction may occur when herbicides are applied under stress (e.g. drought or frost) conditions.

##### **LABELLED ANNUAL AND PERENNIAL BROADLEAVED WEEDS IN CANARYSEED (*Phalaris canariensis*), INCLUDING ANNUAL CANARYGRASS GROWN FOR HUMAN CONSUMPTION**

For the control of labeled annual and perennial broadleaved weeds in Canaryseed (*Phalaris canariensis*), apply Certain Herbicide at a rate of 1.5 – 2.0 L/ha in a minimum of 100 L water/ha to thoroughly cover the weeds with a spray pressure of 200 to 275 kPa. Apply when weeds are in the seedling stage (2 to 4 leaf) and when Canaryseed is in the 3 leaf stage to just before flag leaf emergence stage using a ground boom sprayer. Apply only one post-emergent application per season. Do NOT apply within 60 days of harvest. For use in the Prairie Provinces and the Peace River, Okanagan and Creston regions ONLY of British Columbia. Annual canarygrass seeds can be harvested for human consumption.

REFER TO THE MAIN CERTAIN HERBICIDE LABEL FOR ADDITIONAL DETAILS AND INSTRUCTIONS.

#### **SPRAY BUFFER ZONES**

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop	Spray Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Cereals, flax, grasses, canaryseed ( <i>Phalaris canariensis</i> )	1	1	1	1	4

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray drift buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

### RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Certain Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Certain Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Certain Herbicide or other Group 4 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Sharda Cropchem Limited at 1-844-810-5720

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in any way that is inconsistent with the directions on the label.

All products listed are registered trademarks of their respective companies.