

2020-07-08
2010-3976, 2019-6006

GROUP	2	HERBICIDE
-------	---	-----------

<container label>
[indicates optional text]

DRAFT

DRY FLOWABLE

[FOR USE ON WHEAT (SPRING, WINTER OR DURUM), BARLEY, OATS AND SULFONYLUREA (SU) TOLERANT
CANOLA]

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Thifensulfuron methyl 50% Tribenuron methyl 25%

WARNING – EYE AND SKIN IRRITANT

REGISTRATION. NO.: 31904 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 20 grams – 13.6 kg

Rotam LTD
Unit 6, 26/F Trend Centre,
29 Cheung Lee Street
Chai Wan, Hong Kong

Manufactured for:
Rotam North America, Inc.
330 St. Mary Avenue, Suite 300,
Winnipeg, MB, R3C 3Z5
1-866-927-6826

- **KEEP OUT OF REACH OF CHILDREN.**
- Avoid breathing spray mist. Use with adequate ventilation.
- Avoid contact with skin, eyes and clothing.
- Do not contaminate any body of water.
- May be harmful if swallowed, inhaled or absorbed through skin.
- Causes eye and skin irritation.
- When using, do not eat, drink or smoke. Wash skin thoroughly with soap and water after handling.
- Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before re-use.
- Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves and socks and chemical-resistant footwear during mixing, loading, application, clean-up and repair. In addition, wear a face shield during mixing and loading. Chemical-resistant gloves are not required while operating groundboom sprayers.

FIRST AID:

If in eyes:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or the product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call CHEMTREC 1-800-424-9300 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

If this is a refillable container:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

If this is a returnable container:

Do not reuse this container for any other purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

<booklet>

GROUP	2	HERBICIDE
-------	---	-----------

DRAFT

DRY FLOWABLE

FOR USE ON WHEAT (SPRING, WINTER OR DURUM), BARLEY, OATS AND SULFONYLUREA (SU)
TOLERANT CANOLA

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: Thifensulfuron methyl 50% Tribenuron methyl 25%

WARNING – EYE AND SKIN IRRITANT

REGISTRATION. NO.: 31904 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 20 grams – 13.6 kg

Rotam LTD
Unit 6, 26/F Trend Centre,
29 Cheung Lee Street
Chai Wan, Hong Kong

Manufactured for:
Rotam North America, Inc.
330 St. Mary Avenue, Suite 300,
Winnipeg, MB, R3C 3Z5
1-866-927-6826

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Avoid breathing spray mist. Use with adequate ventilation.
- Avoid contact with skin, eyes and clothing.
- Do not contaminate any body of water.
- May be harmful if swallowed, inhaled or absorbed through skin.
- Causes eye and skin irritation.
- When using, do not eat, drink or smoke. Wash skin thoroughly with soap and water after handling.
- Remove contaminated clothing immediately after use. Store and wash contaminated clothing separately from household laundry before re-use.
- Do not apply by air.
- Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves and socks and chemical-resistant footwear during mixing, loading, application, clean-up and repair. In addition, wear a face shield during mixing and loading. Chemical-resistant gloves are not required while operating groundboom sprayers.
- **Do not** enter or allow entry into treated areas for 12 hours after application.

FIRST AID:**If in eyes:**

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or the product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call CHEMTREC 1-800-424-9300 (24 hours).

TOXICOLOGICAL INFORMATION: Treat symptomatically.

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, recreational grounds or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on other crops.

DO NOT apply using aerial application equipment.

Carefully observe sprayer clean-up instructions, as spray tank residues may damage crops other than wheat, barley, oats or SU Canola™.

APPLY WITH A SURFACTANT REGISTERED FOR USE WITH DRAFT

**GENER
AL
INFOR
MATIO
N**

DRAFT is recommended for selective postemergence control or suppression of broadleaf weeds in wheat (spring, winter or durum), barley, oats and SU tolerant canola. DRAFT is a 75% active ingredient herbicide formulated as a dry flowable dispersible granule. DRAFT is to be mixed with water and a recommended surfactant and applied as a uniform broadcast spray. It is noncorrosive, nonflammable, nonvolatile and does not freeze.

DRAFT MUST be applied with a recommended surfactant, early postemergence, to the main flush of actively growing broadleaf weeds. Recommended surfactants are AGRAL* 90, AG-SURF*, and CITOWETT® PLUS.

DRAFT is a Group 2 herbicide and may only suppress or not control Group 2 resistant weeds.

Warm, moist growing conditions promote active weed growth and enhance the activity of DRAFT by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur. APPLY DRAFT WITH GROUND EQUIPMENT ONLY. For best results, ensure thorough spray coverage of target weeds. See "APPLICATION INSTRUCTIONS" section of this label for complete use details.

DRAFT stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

The following crops may be planted two months after an application of DRAFT: canola, flax, lentils and alfalfa.

**DIRE
CTIO
NS
FOR
USE**

- DO NOT re-enter treated fields until 12 hours after application.
- Apply only when the potential for drift to areas of human habitation or areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

Field sprayer application:

- **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty.
- **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Buffer zones:

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter

belts, woodlots, hedgerows, pastures, rangelands, riparian areas, and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Buffer zone (metres) required for the protection of:		
	Terrestrial habitat	Aquatic habitats of water depths:	
		<1m	>1m
Field sprayer	15	1	0

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

ENVIRONMENTAL PRECAUTIONS:

- **TOXIC** to plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g. soils that are compacted or fine textured such as clay).
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body

SULFONYLUREA TOLERANT CANOLA

(FOR USE ONLY ON SULFONYLUREA TOLERANT CANOLA THAT CONTAINS THE CIBUS SU CANOLA™ TRAIT)

APPLICATION TIMING

Apply 20 grams of DRAFT per hectare with a surfactant which is recommended for use with DRAFT at 2.0 litres per 1,000 litres of spray volume (see "Mixing Instructions" section of this label) when SU Canola™ is at the 2 to 5 leaf stage of development but prior to the beginning of bolting for the control or suppression of weeds listed below. For best results apply to emerged, young, actively growing weeds that are less than 10 cm tall or across. Thorough coverage of target weeds is essential.

RESTRICTIONS:

- Only for use on canola that contains the Cibus sulfonylurea herbicide tolerant trait (SU Canola™ Trait).
- Do not apply to non-sulfonylurea tolerant canola as severe crop injury or death of the plants may occur.
- Do not exceed more than 20 g of product per hectare applied to any one crop of SU tolerant Canola (Cibus SU Canola™) during one growing season.

WEEDS CONTROLLED:

Annual smartweed (green smartweed, lady's thumb)	Narrow-leaved hawk's-beard
Ball mustard	Redroot pigweed
Chickweed ²	Russian thistle
Corn spurry	Shepherd's purse
Cow cockle	Stinkweed
Flixweed	Tartary buckwheat
Groundsel, common	Volunteer rapeseed ⁶

Hemp-nettle
Kochia
Lamb's-quarters

Volunteer sunflower
Wild buckwheat³
Wild mustard

WEEDS SUPPRESSED:

Canada thistle¹
Cleavers (1 to 3 whorls)
Round-leaved mallow⁴
Scentless chamomile
Sow thistle¹
Stork's-bill (2-6 leaf)
Toadflax⁵

Weed suppression is a visual reduction in weed competition (reduced population and/or vigour) as compared to an untreated check. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

Specific Weed Recommendations:

- ¹ Canada thistle, Sow thistle: Apply DRAFT plus recommended surfactant postemergence when the majority of the thistles have emerged and are actively growing. For best top growth control, apply when thistles are no larger than 15 cm tall, before bud stage, and before crop canopy prevents thorough coverage of thistles. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.
- ² Chickweed: Apply DRAFT plus recommended surfactant when the chickweed is small (1-6 leaf) and actively growing but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.
- ³ Wild Buckwheat: Apply DRAFT plus surfactant to actively growing wild buckwheat in the cotyledon - 3 leaf stage. Under stress conditions such as dry weather control may be reduced. Large plants may regrow after treatment.
- ⁴ Round-leaved Mallow: Apply DRAFT plus a surfactant to actively growing round leaved mallow in the 2-6 leaf stage (10-12 cm in height). DRAFT will keep mallow stunted but may not reduce the overall populations.
- ⁵ Toadflax: Apply DRAFT plus recommended surfactant when toadflax is no taller than 15 cm in height. A control program for this weed includes both frequent tillage and chemical application.
- ⁶ Volunteer rapeseed: DRAFT alone will not control imazamox and imazethapyr tolerant canola, (e.g. Clearfield* canola)

NOTE: Avoid making applications to weeds when rainfall is threatening. Rainfall within 4 hours will result in reduced weed control of DRAFT.

Tank Mixtures

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Rotam LTD at 1-866-927-6826 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

DRAFT + LONTREL* 360 or CT MIX 360

For Use on Sulfonylurea Tolerant Canola (for use only on sulfonylurea tolerant canola that contains the Cibus SU Canola™ trait),

APPLY:

DRAFT	at 20 grams per hectare
plus	
LONTREL* 360 or CT MIX 360	at 0.21 litres per hectare
plus	
a recommended surfactant	at 2 litres per 1000 litres of spray volume

See "Mixing Instructions" section of the label for proper mixing/application instructions. Ensure that DRAFT is completely in suspension in the spray tank before adding LONTREL* 360 or CT MIX 360.

Apply the tankmix from the full 2-leaf stage to the 5-leaf stage of SU tolerant canola Consult the LONTREL* 360 or CT MIX 360 label for additional application instructions and use precautions.

WEEDS CONTROLLED:

Canada Thistle (seasonal control) †
Wild Buckwheat ††

PLUS: All other broadleaf weeds controlled or suppressed by DRAFT alone at 20 grams/ha.

†Canada Thistle (seasonal control): Apply DRAFT plus LONTREL* 360 or CT MIX 360 plus a recommended surfactant postemergent, after all the thistles have emerged, and are actively growing. For best control, apply when thistles are no larger than 15 cm tall, before bud stage and before crop canopy prevents thorough coverage of thistles. Later emerging thistles will not be controlled.

†† Wild Buckwheat: Apply DRAFT plus LONTREL* 360 or CT MIX 360 plus a recommended surfactant to actively growing wild buckwheat in the cotyledon to 5-leaf stage. Under stress conditions such as dry weather, control may be reduced. Large plants may regrow after treatment.

DRAFT HERBICIDE + YUMA GL LIQUID EC HERBICIDE

For use on Sulfonylurea Tolerant Canola (for use only on sulfonylurea canola that contains the Cibus SU Canola™ trait)

For the control of annual grasses as well as broadleaf weeds, apply a tank mixture of:

DRAFT HERBICIDE at 20 grams per hectare
plus
YUMA GL LIQUID EC HERBICIDE at 0.38 - 0.5 litres per hectare
plus
MERGE at 0.5 % v/v or 5 litres per 1000 litres spray solution

Apply DRAFT HERBICIDE plus YUMA GL LIQUID EC HERBICIDE tank mixture when the sulfonylurea tolerant canola (Cibus SU Canola™ trait) is in the 3- to 5-leaf stage of growth but prior to beginning of bolting. Application of the DRAFT HERBICIDE plus YUMA GL LIQUID EC HERBICIDE tank mixture prior to the 3-leaf stage of SU tolerant canola could result in crop injury.

This tankmix will control the above broadleaf weeds as well as the grass weeds listed below:

RATE	WEEDS CONTROLLED	WEED LEAF STAGE
0.38 or 0.5 L/ha	Barnyard grass	2 leaf to early tillering*
	Green foxtail	2 leaf to early tillering

Volunteer barley	2 leaf to early tillering
Volunteer corn	2-6 leaf
Volunteer oats	2 leaf to early tillering**
Volunteer wheat	2 leaf to early tillering
Wild oats	1-5 leaf without tillers (0.38 L/ha) 1-5 leaf to early tillering (up to 2 tillers)** (0.5 L/ha)
WEEDS SUPPRESSED	WEED LEAF STAGE
Quackgrass	2-6 leaf*

* Use the higher rate of YUMA GL LIQUID EC HERBICIDE (0.5 L/ha) for control of barnyard grass, as well as suppression of quackgrass.

** Best results on volunteer/wild oats if application is made before tillering begins.

NOTE:

Consult the YUMA GL LIQUID EC HERBICIDE label for additional instructions regarding grassy weed size and/or height at time of application as well as other use precautions, and restrictions.

DRAFT HERBICIDE + ARROW 240 EC HERBICIDE

For use on Sulfonylurea Tolerant Canola (for use only on sulfonylurea tolerant canola that contains the Cibus SU Canola™ trait)

For the control of annual grasses as well as broadleaf weeds, apply a tank mixture of:
DRAFT HERBICIDE at 20 grams per hectare

ARROW 240 EC HERBICIDE plus at 0.19 litres per hectare

ADAMA X-ACT plus at 0.5 % v/v or 5 litres per 1000 litres spray solution

Apply DRAFT HERBICIDE plus ARROW 240 EC tank mixture when the sulfonylurea tolerant canola (Cibus SU Canola™ trait) is in the 3- to 5-leaf stage of growth but prior to beginning of bolting. Application of the DRAFT HERBICIDE plus ARROW 240 EC tank mixture prior to the 3-leaf stage of SU tolerant canola could result in crop injury.

This tankmix will control the above broadleaf weeds as well as the grass weeds listed below:

Grass Species	Leaf Stage	Application Rate of ARROW 240 EC	Application Rate of X-ACT Adjuvant
---------------	------------	----------------------------------	------------------------------------

Wild Oats Volunteer cereals (wheat, barley, oats) Foxtail (green, yellow) Persian darnel Crabgrass (smooth and large) Proso millet Witchgrass Fall panicum Barnyard grass Volunteer corn Volunteer canary grass Quackgrass suppression	2 to 6	0.19 L/ha	0.5% v/v
--	--------	-----------	----------

NOTE:

Consult the ARROW 240 EC label for additional instructions regarding grassy weed size and/or height at time of application as well as other use precautions, and restrictions.

DRAFT Alone

Wheat (Spring, winter and durum), Barley and Oats

Apply 20 grams of DRAFT per hectare with a surfactant which is recommended for use with DRAFT at 2.0 litres per 1,000 litres of spray volume (see "Mixing Instructions" section of this label). Apply postemergence when the crop is in the two-leaf to flag leaf (shot blade) stage of growth, (prior to head emergence) but before the crop canopy will prevent thorough coverage of target weeds. For best results apply to emerged, young, actively growing weeds that are less than 10 cm tall or across. Thorough coverage of target weeds is essential.

WEEDS CONTROLLED:

- | | |
|--|---------------------------------|
| Annual smartweed (green smartweed, lady's thumb) | Narrow-leaved hawk's-beard |
| Ball mustard | Redroot pigweed |
| Chickweed ² | Russian thistle |
| Corn spurry | Shepherd's purse |
| Chickweed | Stinkweed |
| Canada thistle ¹ | Tartary buckwheat |
| Groundsle (common) | Volunteer rapeseed ⁶ |
| Henbane | Volunteer sunflower |
| Round- | Wild buckwheat ³ |
| leaf- | Wild mustard |
| quarters | |
| mallow ⁴ | |
| Scentless chamomile | |
| Sow thistle ¹ | |
| Stork's-bill | |
| (2-6 leaf) | |
| Toadflax ⁵ | |

Weed suppression is a visual reduction in weed competition (reduced population and/or vigour) as compared to an untreated check. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

Specific Weed Recommendations:

^{1.} Canada thistle, Sow thistle: Apply DRAFT plus recommended surfactant postemergence when the majority of the thistles have emerged and are actively growing. For best top growth control, apply

when thistles are no larger than 15 cm tall, before bud stage, and before crop canopy prevents thorough coverage of thistles. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.

2. Chickweed: Apply DRAFT plus recommended surfactant when the chickweed is small (1-6 leaf) and actively growing but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.
3. Wild Buckwheat: Apply DRAFT plus surfactant to actively growing wild buckwheat in the cotyledon - 3 leaf stage. Under stress conditions such as dry weather control may be reduced. Large plants may regrow after treatment.
4. Round-leaved Mallow: Apply DRAFT plus a surfactant to actively growing round leaved mallow in the 2-6 leaf stage (10-12 cm in height). DRAFT will keep mallow stunted but may not reduce the overall populations.
5. Toadflax: Apply DRAFT plus recommended surfactant when toadflax is no taller than 15 cm in height. A control program for this weed includes both frequent tillage and chemical application.
6. Volunteer rapeseed: DRAFT alone will not control imazamox and imazethapyr tolerant canola, (e.g. Clearfield* canola)

NOTE:

Avoid making applications to weeds when rainfall is threatening. Rainfall within 4 hours will result in reduced weed control of DRAFT.

TANK MIXES WITH DRAFT

DRAFT PLUS MCPA

Wheat (Spring, winter and durum), Barley and Oats

Apply DRAFT at 20 grams per hectare plus MCPA Amine or Ester 500 at 0.7 - 1.1 litres per hectare (or acid equivalent to deliver 350 - 560 grams active ingredient per hectare) plus a recommended surfactant at 2 litres per 1,000 litres of spray volume (see "Mixing Instructions" section of the label). Apply the tank mix from the full 3-leaf stage to just before the flag leaf stage of wheat (spring, durum, winter), barley and oats. This tank mix will control:

WEEDS CONTROLLED:

Annual smartweed †	Prickly lettuce
Ball mustard	Ragweed ††
Burdocks (seedling)	Redroot pigweed
Chickweed	Russian pigweed ††
Cocklebur ††	Russian thistle
Corn spurry	Shepherd's purse
Cow cockle	Stinkweed
False ragweed ††	Tartary buckwheat
Flixweed	Tumble mustard
Giant ragweed ††	Volunteer rapeseed †††
Hare's-ear mustard	Wild buckwheat
Hempnettle	Plaintain (common)
Indian mustard	Wild mustard
Kochia ††	Wild radish
Lamb's-quarters mustard	Wormseed

† Green smartweed and lady's thumb

†† Use a minimum of 400 grams MCPA active ingredient per hectare.

††† Including imazamox and imazethapyr tolerant canola (e.g., Clearfield* canola).

WEEDS SUPPRESSED:

Cana
da
thistl
e
Round-leaved mallow
Scentless chamomile
Sow thistle
Stork's-bill (2-6 leaf) (MCPA Ester only)

Follow "Specific Weed Recommendations" for Canada and sow thistle, wild buckwheat, round-leaved mallow and chickweed.

WEEDS CONTROLLED BY DRAFT PLUS MCPA (0.56 L/HA)

Weeds controlled or suppressed by DRAFT alone plus volunteer imazamox and imazethapyr tolerant canola (e.g., Clearfield* canola) (2-4 leaf stage).

Consult the MCPA label for additional application instructions and use precautions .

DRAFT PLUS 2,4-D

Wheat or Barley –Do Not Use On Oats

Apply DRAFT at 20 grams per hectare tank mixed with 2,4-D and a recommended surfactant. Rates and formulation of 2,4-D include:

1. 2,4-D Amine 500 at 0.84 - 1.1 litres per hectare or
2. 2,4-D Ester LV700 at 0.6 - 0.8 litres per hectare or
3. 2,4-D Ester LV600 at 0.7 - 0.9 litres per hectare or
4. Acid equivalent.

A recommended surfactant at 2 litres per 1,000 litres of spray volume must be added (see "Mixing Instructions" section of the label). Apply the tank mix from the full 3-leaf stage to just before the flag leaf stage of wheat (spring, durum, winter) and barley. DO NOT USE ON OATS.

WEEDS CONTROLLED:

Annual smartweed [†]	Prickly lettuce
Ball mustard	Redroot pigweed
Burdocks (seedling)	Russian pigweed
Chickweed	Russian thistle
Cocklebur	Shepherd's purse
Common ragweed	Stinkweed
Corn spurry	Stork's-bill
Cow cockle	(2-6 leaf)
False ragweed	(2,4-D ester only)
Flixweed	Sweet clover
Giant ragweed	Tartary buckwheat
Hare's-ear mustard	Tumble mustard
Hemp-nettle	Volunteer rapeseed ^{††}
Indian mustard	Wild buckwheat
Kochia	Wild mustard
Lamb's-quarters	Wild radish
Plantain (common)	Wormseed mustard

[†] green smartweed and lady's thumb

^{††} including imazamox and imazethapyr tolerant canola varieties (e.g., Clearfield* canola). Consult the 2,4-D label for appropriate use rates.

WEEDS SUPPRESSED:

Canada thistle
Round-leaved mallow
Scentless chamomile
Sow thistle
Toadflax (2,4-D Ester only)

Follow "Specific Weed Recommendations" for Canada and sow thistle, chickweed, wild buckwheat and round-leaved mallow.

Consult the 2,4-D label for additional application instructions and use precautions.

DRAFT PLUS ASSERT* 300 Herbicide

**Spring Wheat and Spring Barley Only
For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only.**

Apply DRAFT at 20 grams per hectare plus ASSERT* 300 Herbicide at 1.3 to 1.6 litres of product per hectare for the control of broadleaf weeds controlled by DRAFT, and wild oats.

Use the low rate of ASSERT* 300 Herbicide when the majority of wild oats are in the 1 - 3 leaf stage and the higher rate if wild oats are in the 4 leaf stage. Application can be made from the 2-leaf up to and including the 6-leaf stage of the crop.

Consult the ASSERT* 300 Herbicide label for additional application instructions; use precautions and recropping information.

Use the spray preparation within 12 hours or product degradation may occur.

**DRAFT + ASSERT* 300 Herbicide
+ MCPA ESTER 500**

**Spring Wheat and Spring Barley Only
For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only**

Apply DRAFT at 20 grams per hectare plus ASSERT* 300 Herbicide at 1.3 - 1.6 litres of product per hectare plus MCPA Ester 500 at 0.7 - 1.1 litres per hectare (or acid equivalent to deliver 350 - 560 grams active ingredient per hectare) for the control of broadleaf weeds controlled by DRAFT plus MCPA, and wild oats.

Apply at the 4-leaf to 6-leaf stage of the crop.

SOIL ZONES	ASSERT* 300 Herbicide RATE
Black and Grey Wooded	1.6 L/ha
Brown and Dark Brown	1.3 L/ha if the majority of the wild oats are in the 1-3 leaf stage. Apply Assert* 300 Herbicide at the 1.6 L/ha if the wild oats are in the 4-leaf stage.

Consult the ASSERT* 300 Herbicide label for additional application instructions, use precautions and recropping information.

Use the spray preparation within 12 hours or product degradation may occur.

DRAFT + PUMA* SUPER

**Spring Wheat (Including Durum) and Barley
For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only.**

APPLY:

DRAFT	at 20 grams per hectare
Plus	
PUMA* SUPER	at 0.5 or 1.0 Litres per hectare

A non-ionic surfactant is NOT required when DRAFT is tankmixed with PUMA* SUPER.

Apply the DRAFT + PUMA* SUPER tankmix to spring and durum wheat at the 1 to 6-leaf plus 3 tillers stage or to barley at the 1 to 5-leaf plus 2 tillers stage. Apply the tankmix to annual grasses in the 1-6 leaf

stage (up to emergence of third tiller). Consult the PUMA* SUPER label for additional information on appropriate weed staging, application instructions, environmental and use precautions.

**WEEDS
CONTROLL
ED:**

Green Foxtail (PUMA* SUPER at 0.5 or 1.0 L/ha)
 Wild Oats (PUMA* SUPER at 1.0 L/ha)
 Barnyard Grass (PUMA* SUPER at 1.0 L/ha)

This tankmix will also control or suppress the broadleaf weeds listed for DRAFT alone at 20 grams/ha.

DRAFT + MCPA ESTER 500 + PUMA* SUPER

**Spring Wheat (Including Durum) and Barley
 For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia
 Only**

APPLY:

DRAFT	at 20 grams per hectare
Plus	
MCPA ESTER 500	at 0.84 Litres per hectare
Plus	
PUMA* SUPER	at 0.5 OR 1.0 Litres per hectare

A non-ionic surfactant is NOT required when DRAFT is tankmixed with MCPA ESTER 500 + PUMA* SUPER. Use ONLY MCPA ESTER 500 in this tankmix and only at the recommended rate or a decrease in wild oat control may be observed.

Apply the DRAFT + MCPA ESTER 500 + PUMA* SUPER tankmix to spring and durum wheat at the 1 to 6-leaf plus 3 tillers stage or to barley at the 1 to 5-leaf plus 2 tillers stage. Apply to annual grasses in the 1-6 leaf stage (up to emergence of third tiller). Consult the MCPA ESTER 500 and PUMA* SUPER labels for additional information on appropriate weed staging, application instructions, environmental and use precautions.

WEEDS CONTROLLED:

Green Foxtail (PUMA* SUPER at 0.5 or 1.0 L/ha)
 Wild Oats (PUMA* SUPER at 1.0 L/ha)
 Barnyard Grass (PUMA* SUPER at 1.0 L/ha)

This tankmix will also control or suppress the broadleaf weeds listed for DRAFT at 20 grams/ha + MCPA ESTER 500 at 0.84 Litres/ha.

DRAFT + PUMA* 120 SUPER

**Spring Wheat (Including Durum) and Barley
 For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia
 Only**

APPLY:

DRAFT	at 20 grams per hectare
Plus	
PUMA* 120 SUPER	at 770 mL/Ha or 385 mL/Ha

A non-ionic surfactant is NOT required when DRAFT is tank mixed with PUMA* 120 SUPER.

Apply the DRAFT + PUMA* 120 SUPER tankmix to spring and durum wheat at the 1 to 6-leaf plus 3 tillers stage or to barley at the 1 to 5-leaf plus 2 tillers stage. Apply the tankmix to annual grasses in the 1-6 leaf stage (up to emergence of third tiller). Consult the PUMA* 120 SUPER label for additional information on appropriate weed staging, application instructions, environmental and use precautions.

WEEDS CONTROLLED:

Green Foxtail (PUMA* 120 SUPER at 385 mL/ha)
 Wild Oats, Green Foxtail, (PUMA* 120 SUPER at 770 mL/ha)
 Yellow Foxtail, Barnyardgrass

Broadleaf weeds controlled or suppressed by DRAFT alone.

DRAFT + PUMA* 120 SUPER + MCPA Ester 500

**Spring Wheat (Including Durum) and Barley
 For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only**

APPLY:

DRAFT	at 20 grams per hectare
Plus	
PUMA* 120 SUPER	at 770 mL/Ha or 385 mL/Ha
Plus	
MCPA Ester 500	0.84 or 0.56 L/Ha

A non-ionic surfactant is NOT required when DRAFT is tank mixed with PUMA* 120 SUPER and MCPA Ester 500.

Use ONLY MCPA ESTER 500 in this tank mix and only at the recommended rate or a decrease in wild oat control may be observed.

Apply the DRAFT + PUMA* 120 SUPER + MCPA Ester 500 tank mix to spring and durum wheat at the 1 to 6-leaf plus 3 tillers stage or to barley at the 1 to 5-leaf plus 2 tillers stage. Apply the tank mix to annual grasses in the 1-6 leaf stage (up to emergence of third tiller). Consult the MCPA Ester 500 and PUMA* 120 SUPER label for additional information on appropriate weed staging, application instructions, environmental and use precautions.

WEEDS CONTROLLED:

Green Foxtail (PUMA* 120 SUPER at 385 mL/Ha)

Wild Oats, Green Foxtail, (PUMA* 120 SUPER at 770 mL/ha)
 Yellow Foxtail, Barnyard
 Grass
 Broadleaf weeds controlled or suppressed by DRAFT alone

Broadleaf weeds controlled by MCPA Ester 500 at 0.84 L/Ha,
OR
 Volunteer imazamox and imazethapyr tolerant canola (e.g., Clearfield*
 canola) (2-4 leaf stage) when MCPA Ester 500 is applied at 0.56 L/ha.

DRAFT + HORIZON* 240 EC

Spring Wheat (Including Durum) Only
For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only

For control of wild oats, apply DRAFT at 20 grams per hectare with HORIZON* 240 EC at 230 mL per hectare and 800 mL per hectare of SCORE* at the 2 leaf to flag leaf stage of the crop. Add DRAFT to the spray tank first, followed by HORIZON* 240 EC, and add SCORE* Adjuvant last. A non-ionic surfactant is NOT required when DRAFT is tank mixed with HORIZON* 240 EC plus the SCORE* adjuvant.

For control of wild oats and green foxtail, apply DRAFT at 20 grams per hectare with HORIZON* 240 EC at 290 mL per hectare and 1 L per hectare of SCORE* at the 2 leaf to flag leaf stage of the crop. Add DRAFT to the spray tank first, followed by HORIZON* 240 EC, and add Score Adjuvant last. A non-ionic surfactant is NOT required when DRAFT is tank mixed with HORIZON* 240 EC plus the SCORE* adjuvant.

Consult the HORIZON* 240 EC label for additional application instructions and use precautions.

WEEDS CONTROLLED:

Wild Oats
 Green Foxtail

This tankmix will also control or suppress the broadleaf weeds listed for DRAFT alone at 20 grams per hectare.

DRAFT + MCPA ESTER 500 + HORIZON* 240 EC

Spring Wheat (Including Durum) Only
For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only

APPLY:

DRAFT	at 20 grams per hectare
Plus	
MCPA Ester 500	at 0.84 - 1.1 L per hectare OR at 0.56 L per hectare
Plus	
HORIZON* 240 EC	at 230 mL per hectare
Plus	
SCORE* Adjuvant	at 0.8% v/v

Apply at the 3 to flag leaf stage of the crop, to actively growing wild oats in the 1 to 6 leaf stage, prior to emergence of the 4th tiller, and to green foxtail in the 1 to 5 leaf stage, prior to emergence of the 3rd tiller. Consult the MCPA Ester 500 and HORIZON* 240 EC labels for additional application instructions and use precautions.

WEEDS

CONTROLLED:

Green Foxtail†

Wild Oats

PLUS broadleaf weeds controlled by DRAFT and broadleaf weeds controlled by MCPA Ester 500 at 0.84-1.1 L per hectare or volunteer imazamox and imazethapyr tolerant canola (e.g. Clearfield* canola) (2-4 leaf stage) when MCPA Ester 500 is applied at 0.56 L per hectare.

†NOTE:

A reduction in control of green foxtail may be observed with the tankmix of DRAFT + HORIZON* + MCPA Ester 500

DRAFT + ACCORD* DF Herbicide + HORIZON* 240 EC

**Spring Wheat (Including Durum) Only
For Use in the Prairie Provinces and Interior (including Peace River Region) of
British Columbia Only**

A
P
P
L
Y
:

DRAFT	at 20 grams per hectare
Plus	
ACCORD* DF Herbicide	at 67 grams per hectare
Plus	
HORIZON* 240 EC	at 230 mL per hectare
Plus	
SCORE* Adjuvant	at 0.8% v/v

Make application when the crop is in the 2-5 leaf stage, to actively growing wild oats in the 1 to 6 leaf stage, prior to emergence of the 4th tiller, and to green foxtail in the 1 to 5 leaf stage, prior to emergence of the 3rd tiller. Thorough coverage of foliage is important for consistent weed control. Prior to use, consult the ACCORD* DF Herbicide label for additional rotational cropping restrictions, application instructions, use precautions and other restrictions. Do not graze the treated crops or cut for hay within 77 days.

WEEDS CONTROLLED:

Green foxtail
Wild oats
Cleavers (1-3 whorl)
PLUS broadleaf weeds controlled or suppressed by DRAFT

DRAFT + LONTREL* 360 or CT MIX 360 + 2,4-D ESTER or MCPA ESTER

**Spring Wheat (Excluding Durum) and Barley Only.
For Use in Western Canada Only.**

APPLY:

DRAFT	at 20 grams per hectare
plus	
LONTREL* 360 or CT MIX 360	at 0.21 litres per hectare
plus	
MCPA Ester (500 g/L)	at 0.84 litres per hectare

or	
2,4-D Ester (600 g/L)	at 0.7 litres per hectare

Add a recommended non-ionic surfactant such as AGRAL* 90, AG-SURF* or CITOWETT® PLUS at a rate of 0.2% of the final spray volume (i.e., 2.0 litres per 1000 litres of spray solution).

See "Mixing Instructions" section of the label.

WEEDS CONTROLLED:

Can
ada
thist
le[†]
Lad
y's
thu
mb
Per
enni
al
sow
thist
le
S
t
i
n
k
w
e
e
d

V
o
l
u
n
t
e
e
r

c
a
n
o
l
a

W
i
l
d

b
u
c
k
w
h
e
a
t

W
i
l
d

m
u
s
t
a
r
d

† Provides seasonal control of Canada thistle

WEEDS SUPPRESSED:

Cleavers

PLUS: all other broadleaf weeds controlled or suppressed by DRAFT alone at 20 grams per hectare. Refer to the MCPA Ester or 2,4-D Ester labels for additional broadleaf weeds controlled at the rates recommended above. Consult the LONTREL* or CT MIX 360 Herbicide and 2,4-D Ester or MCPA Ester labels for additional application instructions, use precautions, and other label restrictions.

DO NOT USE ON OATS.

DRAFT + CURTAIL* M

**Spring Wheat (excluding Durum) and Spring Barley Only.
For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only.**

APPLY:

DRAFT	20 grams per hectare
plus	
CURTAIL* M	1.5 litres per hectare
plus	
a recommended surfactant	at 2 litres per 1,000 litres of spray volume

See "Mixing Instructions" section of the label for proper mixing/application instructions. Ensure that DRAFT is completely in suspension in the spray tank before adding CURTAIL*M.

Apply the tank mix from the full 3-leaf stage of spring wheat and spring barley. Consult the CURTAIL* M label for additional application instructions and use precautions.

WEEDS CONTROLLED:

Canada thistle[†]
Wild buckwheat^{††}

PLUS all other broadleaf weeds controlled or suppressed by DRAFT alone at 20 grams per hectare.

[†] Canada thistle: Apply DRAFT plus CURTAIL* M plus a recommended surfactant postemergence after all the thistles have emerged, and are actively growing. For best control, apply when thistles are no larger than 15 cm tall, before bud stage and before canopy prevents thorough coverage of thistles. Later emerging thistles will not be controlled.

^{††} Wild buckwheat: Apply DRAFT plus CURTAIL* M plus a recommended surfactant to actively growing wild buckwheat in the cotyledon to 5 leaf stage. Under stress conditions such as dry weather, control may be reduced. Large plants may regrow after treatment.

DRAFT + ACCORD* DF HERBICIDE

**SPRING WHEAT (including Durum) AND SPRING BARLEY (grown for feed purposes only). Not Underseeded to Forages.
For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only**

APPLY:

DRAFT	at 20 grams per hectare
plus	
ACCORD* DF Herbicide	At 67 OR at 135-165 grams per hectare
plus	
MERGE* adjuvant	at 1.0% v/v (10 litres per 1000 litres spray volume)

A non-ionic surfactant is NOT required when DRAFT is tank mixed with ACCORD* DF Herbicide plus the MERGE* adjuvant.

Make application when crop is in the 2-5 leaf stage. Use the 165 grams per hectare rate of ACCORD* DF Herbicide when green foxtail is the target weed. Thorough coverage of foliage is important for consistent weed control. Do not apply this tank mixture more often than once every two years. Prior to use, consult the ACCORD* DF Herbicide label for additional rotational cropping restrictions, application instructions, use precautions and other restrictions. Do not graze the treated crops or cut for hay within 77 days of application.

WEEDS CONTROLLED WHEN ACCORD* DF HERBICIDE IS USED AT 135-165 GRAMS/HA	APPLICATION TIMING
Green Foxtail	1-5 leaf, 2 tillers
Barnyard Grass	1-5 leaf
Volunteer Flax	1-8 cm
Cleavers	1-4 whorls
Annual Sowthistle	1-4 leaf
PLUS: Weeds controlled or suppressed by DRAFT alone	

Refer to the ACCORD* DF Herbicide label for additional information on appropriate weed staging at application.

Do not graze the treated crops or cut for hay within 77 days (spring wheat, durum) or 80 days (spring barley) of application.

WEEDS CONTROLLED WHEN ACCORD* DF Herbicide IS USED AT 67 GRAMS/HA	APPLICATION TIMING
Cleavers	1-3 whorls
Annual Sowthistle	1-4 leaf
PLUS: Weeds controlled or suppressed by DRAFT alone	

**DRAFT + ACCORD* DF HERBICIDE + PUMA*
120 SUPER**

SPRING WHEAT (Including Durum) AND SPRING BARLEY (grown for feed purposes only). Not Underseeded to Forages.

For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only

Apply

DRAFT	20 grams per hectare
Accord* DF Herbicide	67 grams per hectare
Puma* 120 Super	0.77 Litres per hectare

Merge* adjuvant is NOT required when DRAFT and Accord* DF Herbicide are tank mixed with Puma* 120 Super.

Apply the DRAFT + Accord* DF Herbicide + Puma* 120 Super tank mix to spring and durum wheat at

the 2 to 5-leaf stage or to barley at the 2 to 4-leaf stage. Apply the tank mix to annual grasses in the 1-6 leaf stage (up to emergence of third tiller), or to cleavers in the 1-4 whorl. Consult the Accord* DF Herbicide and Puma* 120 Super labels for additional information on appropriate weed staging, application instructions, environmental and use precautions.

WEEDS CONTROLLED:

Weeds controlled or suppressed by DRAFT alone
Green Foxtail
Wild Oats
Barnyard Grass
Cleavers

Do not graze the treated crops or cut for hay within 77 days (spring wheat, durum) or 80 days (spring barley) of application.

DRAFT + ATTAIN* HERBICIDE TANK MIX

For Use on Spring Wheat (excluding Durum), and Spring Barley Only. Not Underseeded to Legumes.

For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only

APPLY:

DRAFT	at 20 grams per hectare
plus	
ATTAIN* Herbicide Tank Mix	Attain A (fluroxypyr) at 0.30 L/ha
	Attain B (2,4-D ester) at 0.50 L/ha
plus	
Adjuvants: AGRAL* 90, AG-SURF*, or CITOWETT*	at 2L/1000 litres of spray solution

Apply from the 4-leaf to flag-leaf stage of spring wheat or spring barley and seedling stage to 4-leaf/whorls of weed.

Mixing order is DRAFT, then Attain* A then Attain* B then adjuvant. Refer to MIXING DIRECTIONS section for general instructions. Use a spray volume of 100 L/ha.

Do not permit lactating dairy animals to graze fields within 7 days of application. Do not harvest forage or cut hay within 30 days after application. Withdraw meat animals from treated fields at least 3 days before slaughter. Do not harvest the treated mature crop within 60 days after application. Consult the Attain* Herbicide Tank Mix labels for additional application instructions and use precautions.

WEEDS CONTROLLED:

Weeds controlled or suppressed by DRAFT alone
Cleavers (excluding Group 2 resistant cleavers).

DRAFT + LONTREL* 360 or CT MIX 360

**For Use on Spring Wheat (excluding Durum), and Spring Barley Only
For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only**

APPLY:

DRAFT	at 20 grams per hectare
plus	
LONTREL* 360 or CT MIX 360	at 0.21 litres per hectare
plus	
a recommended surfactant	at 2 litres per 1000 litres of spray volume

See "Mixing Instructions" section of the label for proper mixing/application instructions. Ensure that DRAFT is completely in suspension in the spray tank before adding LONTREL* 360 or CT MIX 360.

Apply the tankmix from the full 3 leaf stage to the flag leaf stage of spring wheat and barley. Consult the LONTREL* 360 or CT MIX 360 label for additional application instructions and use precautions.

WEEDS CONTROLLED:

Canada Thistle (seasonal control) †
Wild Buckwheat ††

PLUS: All other broadleaf weeds controlled or suppressed by DRAFT alone at 20 grams/ha.

†Canada Thistle (seasonal control): Apply DRAFT plus LONTREL* 360 or CT MIX 360 plus a recommended surfactant postemergent, after all the thistles have emerged, and are actively growing. For best control, apply when thistles are no larger than 15 cm tall, before bud stage and before crop canopy prevents thorough coverage of thistles. Later emerging thistles will not be controlled.

†† Wild Buckwheat: Apply DRAFT plus LONTREL* 360 or CT MIX 360 plus a recommended surfactant to actively growing wild buckwheat in the cotyledon to 5-leaf stage. Under stress conditions such as dry weather, control may be reduced. Large plants may regrow after treatment.

DRAFT + BANVEL II* HERBICIDE

For Use on Spring Wheat (excluding Durum), and Spring Barley Only. Not Underseeded to Forages.

For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only

Apply from the 2-5 leaf stage of spring wheat or spring barley and seedling stage to 8 leaf stage of kochia.

DRAFT	20 grams/hectare
Banvel II* Herbicide	0.110 litres/hectare
Adjuvants: Agral*90, Ag-surf* or Citowett* Plus	2 litres/1000 litres of spray solution

Mixing order is DRAFT then Banvel II* Herbicide then adjuvant. Refer to the MIXING DIRECTIONS section for general instructions. Use minimum volume of 55 L/ha.

Do not permit lactating dairy animals to graze fields within 7 days after application. Consult the Banvel II* Herbicide label for additional application instructions and use precautions.

WEEDS CONTROLLED:

Broadleaf weeds controlled or suppressed by DRAFT
Kochia, including Group 2 resistant kochia

DRAFT + HORIZON* 240 EC HERBICIDE + BANVEL II* Herbicide

**For Use on Spring Wheat (excluding Durum). Not Underseeded to Forages.
For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only**

Apply from the 2-5 leaf stage of spring wheat and seedling stage to 8 leaf stage of kochia.

DRAFT	20 grams/hectare
Horizon* 240 EC Herbicide	0.230 L/Ha
Banvel II* Herbicide	0.110 L/Ha
Adjuvant: Score*	0.8 L/Ha

Mixing order is DRAFT then Banvel II* Herbicide then Horizon* 240 EC then Score*. Refer to the MIXING DIRECTIONS section for general instructions. Use minimum volume of 55 L/ha.

Do not permit lactating dairy animals to graze fields within 7 days after application. Consult the Horizon* 240 EC and Banvel II* Herbicide labels for additional application instructions and use precautions.

WEEDS CONTROLLED:

Broadleaf weeds controlled or suppressed by DRAFT
Kochia, including Group 2 resistant kochia
Wild oats
Green foxtail

DRAFT + 2,4-D AMINE OR ESTER + EVEREST 70 WDG

For Use on Spring Wheat (Including Durum) Only

For Use in the Prairie Provinces and Interior (including Peace River Region) of British Columbia Only

Apply one of the following tank mixes from the 1 to 6-leaf (1 to 4 leaves on main stem, plus 2 tillers) growth stage of the crop. For wild oats, green foxtail and volunteer tame oats, apply from the 1 to 6-leaf (1 to 4 leaves on main stem, plus 2 tillers):

TANK MIX COMPONENTS AND APPLICATION RATES								
Component 1		Component 2		Component 3		Component 4		WEEDS CONTROLLED - Broadleaf weeds controlled or suppressed by DRAFT alone plus:
DRAFT	20 g/ha	Everest* 70 WDG	43 g/ha	2,4-D Amine or Ester	up to 560 g ae/ha	Agral * 90 or Agsurf*	0.25% v/v	Wild oats, green foxtail and volunteer tame oats. Use 43 g/ha when wild oat infestations are greater than 100 plants/m ² or when plants are not actively growing due to poor environmental conditions.
DRAFT	20 g/ha	Everest* 70 WDG	28.5 g/ha	2,4-D Amine or Ester	up to 560 g ae/ha	Agral * 90 or Agsurf*	0.25% v/v	Wild oats and volunteer tame oats only. Use 28.5 g/ha when wild oat populations are below 100 plants/m ² and when plants are actively growing.
DRAFT	20 g/ha	Everest* 70 WDG	21.5 g/ha	2,4-D Amine or Ester	up to 560 g ae/ha	Agral * 90 or Agsurf*	0.25% v/v	Green foxtail only

DRAFT	20 g/ha	Everest* 70 WDG	43 g/ha	Agral * 90 or Agsurf*	0.25% v/v			Wild oats, green foxtail, volunteer tame oats. Use 43 g/ha when wild oat infestations are greater than 100 plants/m ² or when plants are not actively growing due to poor environmental conditions.
-------	---------	--------------------	---------	-----------------------------	--------------	--	--	---

Mixing order is Everest 70 WDG then DRAFT then 2,4-D Amine or Ester then adjuvant. Refer to MIXING DIRECTIONS section for general instructions.

Observe minimum interval to harvest of 80 days after treatment. For black, dark brown and gray-wooded soil zones only, field peas may be grown the year following Everest* application in fields where precipitation has been normal or above normal (10 year average) during the growing season, and where organic matter content is above 4% and pH below 7.5. Re- crop spring wheat only in brown soil zone. Consult the 2,4-D Amine or Ester and Everest* 70 WDG labels for additional information on rotational cropping restrictions, application instructions, use precautions and other restrictions.

MIXING INSTRUCTIONS:

1. Select a spray volume that will ensure thorough coverage and uniform spray pattern. Use a minimum of 55-L spray volume per hectare. If tankmixing with other herbicides, always consult the label of the tank mix partner(s) for minimum spray volume requirements and apply the tank mixture using a water volume recommended for all products.
2. Fill the spray tank 1/4 – 1/3 full with fresh water and start agitation.
3. Add the required amount of DRAFT to the tank. Make sure all of the DRAFT is in suspension before proceeding to the next step. Continuous agitation is recommended throughout the entire mixing procedure to ensure products are completely dissolved and/or in suspension.
4. If using DRAFT alone, a recommended surfactant such as AGRAL* 90, AG-SURF* or CITOWETT® PLUS must be added to the spray mixture at a rate of 2 litres per 1000 litres of spray solution.
5. Specific Tank Mix Instructions:
If tankmixing with MCPA, 2,4-D, CURTAIL* M, LONTREL* 360 or CT Mix 360, add the required amount of MCPA, 2,4-D, CURTAIL* M, LONTREL* 360 or CT Mix 360 to the tank while maintaining agitation. Once the herbicide is completely in suspension, add a recommended surfactant such as AGRAL* 90, AG-SURF* or CITOWETT® PLUS at 2 L per 1000 L spray solution.
Anti-foaming agents may be required.

If tankmixing with PUMA* SUPER or MCPA + PUMA* SUPER, add the required amount of PUMA* SUPER and/or MCPA while maintaining agitation. DO NOT ADD A SURFACTANT.

If tank mixing with ASSERT* 300 Herbicide or ASSERT*300 Herbicide + MCPA: once DRAFT is completely in suspension, add the required amount of Spray Water pH Adjuster and agitate until dissolved. Add the required amount of ASSERT* 300 Herbicide while maintaining agitation. If adding MCPA, add it to the tank last. Addition of surfactant is not required with this mixture.

When mixing with LONTREL* 360 + 2,4-D Ester or MCPA Ester, mix DRAFT first in fresh water, then add the required amount of LONTREL* 360 or CT Mix 360 followed by MCPA or 2,4-D. Add the recommended non-ionic surfactant such as AGRAL* 90, AG-SURF* or CITOWETT® PLUS last, at a rate of 2 litres per 1000 litres spray solution.

If tankmixing with HORIZON* 240 EC, or HORIZON* 240 EC + MCPA Ester, add the required amount of HORIZON* 240 EC and/or MCPA Ester after DRAFT is completely in suspension, and maintain agitation. Add the SCORE* adjuvant at the appropriate rate last. DO NOT ADD AN ADDITIONAL NON-IONIC SURFACTANT.

If tankmixing with YUMA GL LIQUID EC HERBICIDE add the required amount of YUMA GL LIQUID EC HERBICIDE after DRAFT is completely in suspension, and maintain agitation. Add the MERGE or AMIGO adjuvant at 0.5 %v/v. DO NOT ADD AN ADDITIONAL NON-IONIC SURFACTANT.

If tankmixing with ARROW 240 EC add the required amount of ARROW 240 EC after DRAFT is completely in suspension, and maintain agitation. Add the ADAMA X-ACT adjuvant at 0.5 %v/v. DO NOT ADD AN ADDITIONAL NON-IONIC SURFACTANT.

When tankmixing with ACCORD* DF Herbicide, add the required amount of DRAFT and ACCORD* to the tank. When these products have completely dissolved and are in suspension, add the MERGE* adjuvant at the appropriate rate last. Do not add an additional non-ionic surfactant.

6. Continue to fill tank and maintain agitation for at least 5 minutes to ensure thorough mixing.
7. On repeat loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of the volume about to be mixed.
8. Use the spray preparation of DRAFT within 24 hours or product degradation may occur. Use spray preparation containing ASSERT* 300 Herbicide within 12 hours or product degradation may occur. If spray solution is left standing without agitation, thoroughly agitate and re-suspend any settled DRAFT before resuming the spraying operation.

The volumetric measuring guide to measure DRAFT is to be used only as a guide. For precise measurement use scales calibrated in grams.

APPLICATION INSTRUCTIONS:

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 55 litres spray volume per hectare). Use 50 mesh filter screens or larger (metal or nylon). A spray pressure of 210-275 kPa is recommended. Continuous agitation is required to keep DRAFT in suspension. Apply uniformly. Avoid overlapping and shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

SPRAYER CLEANUP:

To avoid subsequent injury to crops other than wheat, barley or oats, immediately after spraying and prior to spraying other crops, thoroughly remove all traces of DRAFT from mixing and spray equipment as follows:

1. Drain tank; then flush tank, boom and hoses with clean water for a minimum of ten minutes. Visually inspect tank to ensure removal of all visible residues of DRAFT. If necessary, repeat Step 1. **DO NOT CLEAN SPRAYER NEAR WELL OR WATER SOURCE OR NEAR TO DESIRABLE VEGETATION.**
2. Fill the tank with clean water, then add 1 litre household AMMONIA (containing minimum of 3% ammonia) per 100 litres of water. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation. Again flush the hoses, boom and nozzles with the cleaning solution and drain tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat Steps 2 and 3.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

CAUTION: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which may cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

USE PRECAUTIONS:**RESISTANCE-MANAGEMENT RECOMMENDATIONS:**

For resistance management, DRAFT is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to DRAFT and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the used of DRAFT or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- ◆ Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision

fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

- ◆ Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- ◆ Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local Rotam representative at 1-866-927-6826 for further information.

- ◆ Applications to wheat, barley or oats that are stressed by severe weather conditions, frost, low fertility, drought, water logged soil (soils at or near field capacity), disease or insect damage may result in crop injury. Drought, disease or insect damage following application may also result in crop injury.

- ◆ Do not apply to irrigated land where tail water will be used to irrigate cropland.
- ◆ Do not contaminate irrigation water.
- ◆ Do not apply to crops underseeded to legumes or grasses as injury to forage may result.
- ◆ Do not remove soil from treated fields for use on lawns, in gardens or for back fill.
- ◆ DRAFT should be applied only with water as a carrier. Other carriers may accelerate breakdown of the product and reduce its effectiveness.
- ◆ Do not exceed a total of 20 g of DRAFT per hectare per crop year.
- ◆ Do not plant to any crop until two months after application.
- ◆ Wheat, barley or oats may not be grazed by or fed to livestock within 7 days of application.
- ◆ Under certain conditions (such as heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures) temporary lightening in crop colour and occasionally a slight reduction in crop height may occur.
- ◆ Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls, or calcium deposits.

- ◆ Limit use of DRAFT to currently licensed varieties of wheat (spring, durum, winter), barley, oats and SU tolerant canola. Consult Rotam prior to use on unlicensed varieties for crop tolerance information. **DO NOT USE ON LEGER BARLEY AND BELVEDERE WHEAT.**

Overspray or drift to important wildlife habitats such as shelterbelts, wetlands, sloughs or dry slough borders and woodlots should be avoided.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

If this is a refillable container:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

If this is a returnable container:

Do not reuse this container for any other purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For all package types:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

*All other products mentioned are trademarks of their respective companies