



# Thrasher® II

## Bromoxynil & 2,4-D Ester

Herbicide – Broadleaf Weed Control

Provides excellent dual modes of action for control of 26 tough broadleaf weeds (including Group 2 resistant kochia) in wheat and barley

**Active ingredients:** Bromoxynil and 2,4-D ester

**Chemistry group:** Group 4 (2,4-D) and Group 6 (bromoxynil)

**How it works:** A combination of systemic and contact activity with weeds yellowing within 2 to 4 days and exhibiting abnormal growth (twisting and cupping of leaves) in 2 to 10 days.

**Registered crops:**

- Barley
- Spring wheat (including durum)

**Weeds controlled\*:**

4-leaf stage unless otherwise noted:

- American nightshade
- Ball mustard
- Bluebur
- Cocklebur
- Common buckwheat (8 leaf)
- Common groundsel (8 leaf)
- Common ragweed
- Cow cockle
- Flixweed
- Green smartweed
- Kochia 2 inches high (1–12 leaf)
- Lady's-thumb
- Lamb's-quarters (8 leaf)
- Night-flowering catchfly
- Pale smartweed
- Redroot pigweed
- Russian thistle (2–12 leaf)
- Shepherd's-purse
- Stinkweed (8 leaf)
- Tartary buckwheat (8 leaf)
- Triazine resistant pigweed
- Velvetleaf (3 inches high)
- Volunteer canola
- Volunteer sunflower
- Wild buckwheat (8 leaf)
- Wild mustard (8 leaf)

\*Weeds can be controlled up to the recommended stage

**Crop staging:** 4 leaf to early flag leaf


**Application rates and packaging:**

- 500 mL/ac or 40 ac/case
- 2 x 10L jugs/case

**Water volume:**

- Ground: 20 to 40 L/ac
- Aerial: 12 to 16 L/ac

**Rainfastness:** 1 hour





### Registered and supported tank mixes:

Spring wheat (including durum) and barley:

- Avenge®
- Fenoxaprop (Bengal® WB)
- Tralkoxydim (Bison®)

Spring wheat (including durum) only:

- Clodinafop (Ladder®)
- Everest®

### Mixing instructions:

1. Half fill the tank with clean water
2. Add the required amount of Thrasher and agitate thoroughly
3. Fill the tank and agitate again before use

**Crop rotations:** No restrictions the year after application

**Pre-harvest interval:** 30 days

### Grazing restriction:

- Do not permit livestock to graze fields within 30 days after application
- Do not harvest forage or cut for hay within 30 days after application
- Withdraw meat animals from treated fields at least 3 days before slaughter

### Storage:

- Will solidify at temperatures below -20°C but will become useable at temperatures above 0°C
- Shake well before using

### Quick tips:

- For best results when there is a heavy crop canopy, or when the majority of the weeds are cow cockle, smartweed or pigweed, use higher water volumes
- For best results spray when weeds are in the seedling stage
- Apply in good growing conditions
- Application must be made before the crop shields the weeds

